

The Variational Sequence on Finite Jet Bundle Extensions and the Lagrangian Formalism

D. R. Grigore ¹

Dept. of Theor. Phys., Inst. Atomic Phys.
Bucharest-Măgurele, P. O. Box MG 6, ROMÂNIA

Abstract

The geometric Lagrangian theory (of arbitrary order) is based on the analysis of some basic mathematical objects such as: the contact ideal, the (exact) variational sequence, the existence of Euler-Lagrange and Helmholtz-Sonin forms, etc. In this paper we give new and much simpler proofs for the whole theory using Fock space methods. Using these results we give the most general expression for a variationally trivial Lagrangian and the generic expression for a locally variational system of partial differential equation.

¹e-mail: grigore@theor1.ifa.ro, grigore@roifa.ifa.ro

1 Introduction

The modern Lagrangian theory is formulated in the language of jet bundle extensions. There are two different, but closely related approaches in this framework. One is based on infinite jet bundle extensions [3], [2], [27] (and uses the so-called variational bicomplex) and the other is based on finite jet bundle extensions [17] - [24] (and uses the so-called variational sequence). The second approach is more in the spirit of the original Lagrangian theory as studied in physical literature corresponding to a Lagrangian theory of arbitrary, but finite order. On the other hand, the study of the finite order seems to be much more complicated from the combinatorial point of view as it is apparent from the proofs appearing in the references quoted above. In some recent papers we have tried to prove that the simplest way to disentangle the combinatorial proofs is to use the observation that the various tensors appearing in the line of argument have some symmetry or antisymmetry properties which make them as elements in some Fock space. Then most of the relations to be solved can be written with the help of the creation and annihilation operators. Using elementary properties of these Fock space operators one can significantly simplify the proofs. This is one of the main motivation of writing this paper, i.e to clarify the technical aspects. It appears that it is convenient to write the paper in a self contained style so it will also serve as a pedagogical introduction to this field of interest. In Section 2 we provide the basic construction - the (finite) jet bundle extension idea and give some relevant formulæ. In Sections 3 and 4 we study the contact and respectively the strong contact forms on a jet bundle extension and obtain with our method the structure of the contact ideal. In section 5 we establish the existence of the Euler-Lagrange and of Helmholtz-Sonin forms adapting to the finite jet bundle extension approach some ideas of Anderson and collaborators. The proof of the existence of the Helmholtz-Sonin form is new. In Section 6 we sketch the proof for the exactness of the variational sequence and we clarify some points about the characterization of the elements of the variational sequence by forms. In Section 7 we give the most general expression for a variationally trivial Lagrangian and in Section 8 we provide the generic expression for a locally variational partial differential system of equations using the methods developed in the preceding sections. These problems have been under intensive study for sometime in the literature but many of the results obtained here are new. In particular, we obtain in Section 8 an *if and only if* type of result. Section 9 is devoted to some final comments.

2 Finite Order Jet Bundle Extensions

The content of this section is standard and is included mainly to fix the notations. For the sake of completeness, we give however some proofs which are not trivial.

The kinematical structure of a classical field theory is based on fibred bundle structures. Let $\pi : Y \mapsto X$ be fibre bundle, where X and Y are differentiable manifolds of dimensions $\dim(X) = n$, $\dim(Y) = m + n$ and π is the canonical projection of the fibration. Usually X is interpreted as the “space-time” manifold and the fibres of Y as the field variables. An *adapted chart* to the fibre bundle structure is a couple (V, ψ) where V is an open subset of Y and $\psi : V \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ is the so-called *chart map*, usually written as $\psi = (x^i, y^\sigma)$ ($i = 1, \dots, n; \sigma = 1, \dots, m$) such that $(\pi(V), \phi)$ where $\phi = (x^i)$ ($i = 1, \dots, n$) is a chart on X and the canonical projection has the following expression: $\pi(x^i, y^\sigma) = (x^i)$. If $p \in Y$ then the real numbers $x^i(p)$, $y^\sigma(p)$ are called the (*fibred*) *coordinates* of p . For simplicity we will give up the attribute *adapted* sometimes in the following. Also we will refer frequently to the first entry V of (V, ψ) as a chart.

Next, one considers the *r-jet bundle extensions* $J_n^r Y \mapsto X$ ($r \in \mathbb{N}$). The construction is the following (see for instance [20]).

Theorem 2.1 *Let $x \in X$, and $y \in \pi^{-1}(x)$. We denote by $\Gamma_{(x,y)}$ the set of sections $\gamma : U \rightarrow Y$ such that: (i) U is a neighbourhood of x ; (ii) $\gamma(x) = y$. We define on $\Gamma_{(x,y)}$ the relationship “ $\gamma \sim \delta$ ” iff there exists a chart (V, ψ) on Y such that γ and δ have the same partial derivatives up to order r in the given chart i.e.*

$$\frac{\partial^k}{\partial x^{i_1} \dots \partial x^{i_k}} \psi \circ \gamma \circ \phi^{-1}(\phi(x)) = \frac{\partial^k}{\partial x^{i_1} \dots \partial x^{i_k}} \psi \circ \delta \circ \phi^{-1}(\phi(x)), \quad k \leq r. \quad (2.1)$$

Then this relationship is chart independent and it is an equivalence relation.

Proof: (i) We first prove the chart independence. Let $(V, \psi), (\bar{V}, \bar{\psi})$ be a two (adapted) chart on this bundle such that $V \cap \bar{V} \neq \emptyset$ and let $(\pi(V), \phi), (\pi(\bar{V}), \bar{\phi})$ the corresponding charts on X . If we define $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by:

$$f \equiv \bar{\phi} \circ \phi^{-1}$$

then the function $\bar{\psi} \circ \psi^{-1} : \mathbb{R}^{m+n} \rightarrow \mathbb{R}^{m+n}$ has the following expression:

$$\bar{\psi} \circ \psi^{-1} = (f^i, F^\sigma) \quad (2.2)$$

where F is a smooth function from \mathbb{R}^n into \mathbb{R}^{m+n} . Let $\gamma \in \Gamma_{(x,y)}$; then in adapted coordinates we have the following expression:

$$\psi \circ \gamma \circ \phi^{-1}(x^i) = (x^i, g^\sigma(x)). \quad (2.3)$$

Similarly we have in the other chart:

$$\bar{\psi} \circ \gamma \circ \bar{\phi}^{-1}(x^i) = (x^i, \bar{g}^\sigma(x))$$

where:

$$\bar{g}^\sigma(x) = F^\sigma(f^{-1}(x), g^\sigma(f^{-1}(x))). \quad (2.4)$$

We must show that if g^σ and g'^σ have the same partial derivatives up to order r then \bar{g}^σ and \bar{g}'^σ have the same property. To prove this one firstly gets from the expression of \bar{g}^σ above that:

$$\frac{\partial \bar{g}^\sigma}{\partial x^i} = (\Delta_i F^\sigma)(f^{-1}(x), g^\sigma(f^{-1}(x))) \quad (2.5)$$

where the differential operator Δ_i acts in the space of smooth functions $f : \mathbb{R}^{m+n} \rightarrow \mathbb{R}$ according to:

$$\Delta_i \equiv P_i^j \left(\frac{\partial}{\partial x^j} + \frac{\partial g^\sigma}{\partial x^j} \frac{\partial}{\partial y^\sigma} \right) \quad (2.6)$$

and P_i^j is the inverse of the matrix $\frac{\partial f^l}{\partial x^p}$:

$$P_i^j \frac{\partial f^i}{\partial x^p} = \delta_p^j. \quad (2.7)$$

From (2.5) we obtain by recurrence:

$$\frac{\partial^k \bar{g}^\sigma}{\partial x^{i_1} \dots \partial x^{i_k}} = (\Delta_{i_1} \dots \Delta_{i_k} F^\sigma)(f^{-1}(x), g^\sigma(f^{-1}(x))). \quad (2.8)$$

But it is clear that the expression $\Delta_{i_1} \dots \Delta_{i_k} F^\sigma$ contains the derivatives of the function g^σ only up to order r . So, the chart independence of the relation \sim follows.

(ii) Using the chart independence proved above, it easily follows that \sim is an equivalence relation. ■

A r -order jet with source x and target y is, by definition, the equivalence class of some section γ with respect to the equivalence relationship defined above and it is denoted by $j_x^r \gamma$.

Let us define $J_{(x,y)}^r \pi \equiv \Gamma_{(x,y)} / \sim$ Then the r -order jet bundle extension is, set theoretically $J^r Y \equiv \bigcup_x J_{(x,y)}^r \pi$.

Now we define the following projections: $\pi^{r,s} : J^r Y \rightarrow J^s Y$ ($0 < s \leq r$) by

$$\pi^{r,s}(j_x^r \gamma) = j_x^s \gamma, \quad (2.9)$$

$\pi^{r,0} : J^r Y \rightarrow Y$ given by

$$\pi^{r,0}(j_x^r \gamma) = \gamma(x). \quad (2.10)$$

(this is consistent with the identification $J^0 Y \equiv Y$) and finally $\pi^r : J^r Y \rightarrow X$ by

$$\pi^r(j_x^r \gamma) = x. \quad (2.11)$$

These projections are obviously surjective.

Let (V, ψ) , $\psi = (x^i, y^\sigma)$ be a chart on Y . Then we define the couple (V^r, ψ^r) , where: $V^r = (\pi^{r,0})^{-1}(V)$ and

$$\psi = (x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_k}^\sigma, \dots, y_{j_1, \dots, j_r}^\sigma), \quad j_1 \leq j_2 \leq \dots \leq j_k, \quad k = 1, \dots, r$$

where

$$\begin{aligned} y_{j_1, \dots, j_k}^\sigma(j_x^r \gamma) &= \frac{\partial^k}{\partial x^{j_1} \dots \partial x^{j_k}} y^\sigma \circ \gamma \circ \phi^{-1} \Big|_{\phi(x)}, \quad k = 1, \dots, r \\ x^i(j_x^r \gamma) &= x^i(x), \quad y^\sigma(j_x^r \gamma) = y^\sigma(\gamma(x)). \end{aligned} \quad (2.12)$$

Then (V^r, ψ^r) is a chart on $J^r Y$ called *the associated chart* of (V, ψ) .

Remark 2.1.1 The expressions $y_{j_1, \dots, j_k}^\sigma(j_x^r \gamma)$ are defined for all indices $j_1, \dots, j_k = 1, \dots, n$, and the restrictions $j_1 \leq j_2 \leq \dots \leq j_k$ in the definition of the charts are in order to avoid over-counting and are a result of the obvious symmetry property:

$$y_{j_{P(1)}, \dots, j_{P(k)}}^\sigma(j_x^r \gamma) = y_{j_1, \dots, j_k}^\sigma(j_x^r \gamma), \quad (2.13)$$

for any permutation $P \in \mathcal{P}_k$, $k = 2, \dots, r$.

Now we have the following result.

Theorem 2.2 If a collection of (adapted) charts (V, ψ) are the elements of a differentiable atlas on Y then (V^r, ψ^r) are the elements of a differentiable atlas on $J_n^r(Y)$ which admits a fibre bundle structure over Y .

Proof:

As in theorem 2.1 we consider two non-overlapping charts $(V, \psi), (\bar{V}, \bar{\psi})$ and let $(\pi(V), \phi), (\pi(\bar{V}), \bar{\phi})$ the corresponding charts on X . If we define $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $F : \mathbb{R}^{m+n} \rightarrow \mathbb{R}^m$ as in the proof of this theorem, then we have

$$\bar{\psi}^r \circ (\psi^r)^{-1} = (f^i, F^\sigma, F_j^\sigma, \dots, F_{j_1, \dots, j_r}^\sigma) \quad (2.14)$$

where $F_{j_1, \dots, j_k}^\sigma$, $j_1 \leq j_2 \leq \dots \leq j_k$, $k = 1, \dots, r$ are functions depending on the variables $(x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_k}^\sigma, \dots, y_{j_1, \dots, j_r}^\sigma)$.

We must prove that these functions are smooth. To this purpose we obtain an rather explicit formula for them. Inspecting the notations from theorem 2.1 it is clear that

$$y_{j_1, \dots, j_k}^\sigma = \frac{\partial g^\sigma}{\partial x^{j_1} \dots \partial x^{j_k}} \Big|_{\phi(x)} \quad (2.15)$$

and

$$\bar{y}_{j_1, \dots, j_k}^\sigma = \frac{\partial \bar{g}^\sigma}{\partial x^{j_1} \dots \partial x^{j_k}} \Big|_{\bar{\phi}(x)}.$$

But we have an explicit formula for the last expression, namely (2.8). By induction one proves that the expression $\Delta_{i_1} \dots \Delta_{i_k} F^\sigma$ is a polynomial in the expressions

$$g_{j_1, \dots, j_l}^\sigma = \frac{\partial g^\sigma}{\partial x^{j_1} \dots \partial x^{j_k}} \Big|_{\phi(x)}, \quad l = 1, \dots, k$$

with coefficients depending only on (x^i, y^σ) . But we immediately have

$$\Delta_{j_1} g_{j_2, \dots, j_l}^\sigma = P_{j_1}^i g_{i, j_2, \dots, j_l}^\sigma, \quad l = 1, \dots, k.$$

So one can replace $\Delta_i \rightarrow \tilde{\Delta}_i$ where $\tilde{\Delta}_i$ is an operator acting on polynomials in $g_{j_1, \dots, j_l}^\sigma$, $l = 1, \dots, k$ with coefficients depending only on (x^i, y^σ) according to

$$\tilde{\Delta}_{j_1} g_{j_2, \dots, j_l}^\sigma = P_{j_1}^i g_{i, j_2, \dots, j_l}^\sigma, \quad l = 1, \dots, k.$$

and

$$\tilde{\Delta}_j f = \Delta_j f$$

on the coefficients which are functions depending only on (x^i, y^σ) .

Taking into account (2.15) it follows that $F_{j_1, \dots, j_k}^\sigma$, $k = 1, \dots, r$ is a polynomial in $y_{j_1, \dots, j_l}^\sigma$ for $l = 1, \dots, k$ with smooth coefficients of (x^i, y^σ) so these are smooth functions. The existence of the differentiable structure is proved. The fibre bundle structure over X is obvious. ■

To be able to use the summation convention over the dummy indices we consider $y_{j_1, \dots, j_k}^\sigma$ for *all* values of the indices $j_1, \dots, j_k \in \{1, \dots, n\}$ as smooth functions on the chart V^r defined in terms of the independent variables $y_{j_1, \dots, j_k}^\sigma$, $j_1 \leq j_2 \leq \dots \leq j_k$ $k = 1, 2, \dots, r$ according to the formula (2.13) and we make a similar convention for the partial derivatives $\frac{\partial}{\partial y_{j_1, \dots, j_k}^\sigma}$.

Then we define on the chart V^r the following vector fields:

$$\partial_\sigma^{j_1, \dots, j_k} \equiv \frac{r_1! \dots r_n!}{k!} \frac{\partial}{\partial y_{j_1, \dots, j_k}^\sigma}, \quad k = 1, \dots, r \quad (2.16)$$

for all values of the indices $j_1, \dots, j_k \in \{1, \dots, n\}$. Here r_l , $l = 1, \dots, n$ is the number of times the index l enters into the set $\{j_1, \dots, j_k\}$.

One can easily verify the following formulæ:

$$\partial_\sigma^{i_1, \dots, i_k} y_{j_1, \dots, j_l}^\nu = 0, \quad (k \neq l) \quad (2.17)$$

$$\partial_\sigma^{i_1, \dots, i_k} y_{j_1, \dots, j_k}^\nu = \mathcal{S}_{j_1, \dots, j_k}^+ \delta_{j_1}^{i_1} \dots \delta_{j_k}^{i_k} \quad (2.18)$$

where $\mathcal{S}_{j_1, \dots, j_k}^+$ is the symmetrization projector operator in the indices j_1, \dots, j_k defined by the formula (10.8) from the Appendix.

Also we have for any smooth function f on the chart V^r :

$$df = \frac{\partial f}{\partial x^i} dx^i + \sum_{k=0}^r (\partial_\sigma^{j_1, \dots, j_k} f) dy_{j_1, \dots, j_k}^\sigma = \frac{\partial f}{\partial x^i} dx^i + \sum_{|J| \leq r} (\partial_\sigma^J f) dy_J^\sigma. \quad (2.19)$$

In the last formula we have introduced the multi-index notations in an obvious way. This formula also shows that the coefficients appearing in the definition (2.16) are exactly what is needed to use the summation convention over the dummy indices without overcounting.

We now define the expressions

$$d_i^r \equiv \frac{\partial}{\partial x^i} + \sum_{k=0}^{r-1} y_{i, j_1, \dots, j_k}^\sigma \partial_\sigma^{j_1, \dots, j_k} \quad (2.20)$$

called *formal derivatives*. When it is no danger of confusion we denote simply $d_i = d_i^r$.

Remark 2.2.1 *The formal derivatives are not vector fields on $J^r Y$.*

Next one immediately sees that

$$d_i y_{j_1, \dots, j_k}^\sigma = y_{i, j_1, \dots, j_k}^\sigma, \quad k = 0, \dots, r-1 \quad (2.21)$$

and one can obtain from theorem 2.2 a recurrence relation for the chart transformation formulæ:

Lemma 2.3 *The following recurrence relation are valid for the chart transformation expressions (2.14):*

$$F_{j_1, \dots, j_k}^\sigma = Q_{j_1}^l d_l F_{j_2, \dots, j_k}^\sigma, \quad k = 1, \dots, r-1. \quad (2.22)$$

In particular, the functions $F_{j_1, \dots, j_k}^\sigma$ are polynomial expressions in the variables $y_j^\sigma, \dots, y_{j_1, \dots, j_k}^\sigma$ (with smooth coefficients depending on (x^i, y^σ)) for all $k = 1, \dots, r$.

From the definition of the formal derivatives it easily follows by direct computation that:

$$[\partial_\sigma^{j_1, \dots, j_k}, d_i] = \frac{1}{k} \sum_{l=1}^k \delta_i^{j_l} \partial_\sigma^{j_1, \dots, \hat{j}_l, \dots, j_k}, \quad k = 0, \dots, r \quad (2.23)$$

where we use Bourbaki conventions $\sum_\emptyset \equiv 0$, $\prod_\emptyset \equiv 1$.

Based on this relation one gets:

$$[d_i, d_j] = 0 \quad (2.24)$$

so one can consistently define for any multi-index

$$d_J \equiv \prod_{i \in J} d_i. \quad (2.25)$$

If γ is a section of the fibre bundle Y then the map $X \ni x \mapsto j_x^r \gamma \in J^r Y$ is section of the fibre bundle $J^r Y$ called the *r-extension* of γ . We denote it by: $j^r \gamma : V \rightarrow J^r Y$.

Then one can define a map $h : T(J^{r+1} Y) \rightarrow T(J^r Y)$ called *horizontalization* by

$$h = h_{j_x^{r+1} \gamma} \equiv (j^r \gamma)_{*,x} \circ (\pi^{r+1})_{*,j_x^{r+1} \gamma}. \quad (2.26)$$

If $\xi \in T_{j_x^{r+1} \gamma}(J^{r+1} Y)$ then $h\xi$ is called its *horizontal* component. We will also define

$$p\xi \equiv (\pi^{r+1,r})_* \xi - h\xi. \quad (2.27)$$

Let $\pi_i : Y_i \rightarrow X_i$, $i = 1, 2$ be two fibre bundles. Then a map $\phi : Y_1 \rightarrow Y_2$ is called a *fibre bundle morphism* if there is a map $\phi_0 : X_1 \rightarrow X_2$ such that

$$\pi_2 \circ \phi = \phi_0 \circ \pi_1;$$

one also says that ϕ covers ϕ_0 .

In this case, one can define the *r-order jet extension* of ϕ as the the map $j^r \phi : J^r Y_1 \rightarrow J^r Y_2$ given by

$$j^r \phi(j_x^r \gamma) \equiv j_{\phi_0(x)}^r \phi \circ \gamma \circ \phi_0^{-1}, \quad \forall x \in X_1. \quad (2.28)$$

If ξ is a projectable vector field on the fibre bundle Y i.e. there is a vector field ξ_0 on X such that $\xi_0 = \pi_*\xi$, then the flow ϕ_t^ξ associated to ξ covers the flow $\phi_t^{\xi_0}$ associated to ξ_0 so we can define the *r-order extension* of ξ by

$$j^r\xi \equiv \left. \frac{d}{dt} \right|_{t=0} j^r\phi_t^\xi. \quad (2.29)$$

The vector field $j^r\xi$ is projectable.

A π -vertical vector field on the fibre bundle Y is called a *evolution*. In the chart (V, ψ) an evolution has the expression:

$$\xi = \xi^\sigma \partial_\sigma \quad (2.30)$$

with ξ^σ smooth functions on the chart V . One denotes the set of evolutions by $\mathcal{E}(J^rY)$; this set is a fibre bundle over J^rY .

We denote the forms of degree q on J^rY by Ω_q^r . A form $\rho \in \Omega_q^r$ is called *π^r -horizontal* (or *basic*) if $i_\xi\rho = 0$ for any π^r -vertical vector field on J^rY . In local coordinates such a form has the following expression:

$$\rho = B_{i_1, \dots, i_q} dx^{i_1} \wedge \dots \wedge dx^{i_q} \quad (2.31)$$

with B_{i_1, \dots, i_q} smooth symmetric functions on V^r . We denote the set of basic forms of degree q by $\Omega_{q,X}^r$. The elements of $\lambda \in \Omega_{n,X}^r$ are called *Lagrange forms*. They have the local expression

$$\lambda = L\theta_0 \quad (2.32)$$

where L is a smooth function on V^r and

$$\theta_0 \equiv dx^1 \wedge \dots \wedge dx^n. \quad (2.33)$$

We also define some generalisation of the horizontal forms for $q > n$. We say that $\rho \in \Omega_q^r$ $q > n$ is *horizontal* iff

$$i_{\xi_1} \dots i_{\xi_{q-n+1}} \rho = 0 \quad (2.34)$$

for any vector fields $\xi_1, \dots, \xi_{q-n+1}$ which are π^r -vertical. The local expression of such a form in the chart (V^r, ψ^r) is:

$$\rho = \sum_{|J_1|, \dots, |J_{q-n}| \leq r} A_{\nu_1, \dots, \nu_{q-n}}^{J_1, \dots, J_{q-n}} dy_{I_1}^{\nu_1} \wedge \dots \wedge dy_{I_{q-n}}^{\nu_{q-n}} \wedge \theta_0 \quad (2.35)$$

with $A_{\nu_1, \dots, \nu_{q-n}}^{J_1, \dots, J_{q-n}}$ smooth functions on V^r with appropriate symmetry properties. We conserve the notation $\Omega_{q,X}^r$ for these forms.

3 Contact Forms

In this section we will give a new proof for the structure formula of the contact forms. This problem was solved by different methods in [23] (a sketch of the proof also appears in [21]). Our proof is based on some Fock space methods, as said in the Introduction. These kind of methods have been employed recently for related problems [10], [11]. We start with the basic definitions and some elementary properties.

3.1 Basic Definitions and Properties

By a *contact form* we mean any form $\rho \in \Omega_q^r$ verifying

$$(j^r \gamma)^* \rho = 0 \quad (3.1)$$

for any section γ . We denote by $\Omega_{q(c)}^r$ the set of contact forms of degree $q \leq n$. If one considers only the contact forms on an open set $V \subset Y$ then we emphasize this by writing $\Omega_{q(c)}^r(V)$. One immediately notes that $\Omega_{0(c)}^r = 0$ and that for $q > n$ any q -form is contact. It is also elementary to see that the set of all contact forms is an ideal, denoted by $\mathcal{C}(\Omega^r)$, with respect to the operation \wedge . Because the operations of pull-back and of differentiation are commuting this ideal is left invariant by exterior differentiation:

$$d\mathcal{C}(\Omega^r) \subset \mathcal{C}(\Omega^r). \quad (3.2)$$

By elementary computations one finds out that for any chart (V, ψ) on Y , every element of the set $\Omega_{1(c)}^r(V)$ is a linear combination of the following expressions:

$$\omega_{j_1, \dots, j_k}^\sigma \equiv dy_{j_1, \dots, j_k}^\sigma - y_{i, j_1, \dots, j_k}^\sigma dx^i, \quad k = 0, \dots, r-1 \quad (3.3)$$

or, in multi-index notations

$$\omega_J^\sigma \equiv dy_J^\sigma - y_{iJ}^\sigma dx^i, \quad |J| \leq r-1. \quad (3.4)$$

From the definition above it is clear that the linear subspace of the 1-forms is generated by dx^i , ω_J^σ , ($|J| \leq r-1$) and dy_I^σ , $|I| = r$.

Formula (2.19) can be now written as follows: for any smooth function on V^r we have

$$df = (d_i f) dx^i + \sum_{|J| \leq r-1} (\partial_\sigma^J f) \omega_J^\sigma + \sum_{|I|=r} (\partial_\nu^I f) dy_I^\nu. \quad (3.5)$$

We also have the formula

$$d\omega_J^\sigma = -\omega_{j_i}^\sigma \wedge dx^i, \quad |J| \leq r-2. \quad (3.6)$$

Let us now consider an arbitrary form $\rho \in \Omega_q^r$, $q > 1$. For any $k = 0, \dots, q$ we define its *contact component of order k* to be the form $p_k \rho \in \Omega_q^{r+1}$ given by:

$$p_k \rho(j_x^{r+1} \gamma)(\xi_1, \dots, \xi_q) \equiv \frac{1}{k!(q-k)!} \varepsilon^{j_1, \dots, j_q} \rho(j_x^r \gamma)(p\xi_{j_1}, \dots, p\xi_{j_k}, h\xi_{j_{k+1}}, \dots, h\xi_{j_q}). \quad (3.7)$$

One usually calls $h\rho \equiv p_0\rho$ and $p\rho \equiv \sum_{k=1}^q p_k\rho$ the *horizontal component* and respectively the *contact component* of ρ . It is useful to particularize the definition above for the case $k = 0$:

$$h\rho(\xi_1, \dots, \xi_q) = \rho(h\xi_1, \dots, h\xi_q). \quad (3.8)$$

A form $\rho \in \Omega_q^r$ is called *k-contact* if $p_j\rho = 0, \forall j \neq k$ and one says that it has the *contact order greater than k* if $p_j\rho = 0, \forall j \leq k-1$.

Now we have a decomposition property, namely

Proposition 3.1 *For any $\rho \in \Omega_q^r$ the following formula is valid:*

$$(\pi^{r+1,r})^*\rho = \sum_{k=0}^q p_k\rho. \quad (3.9)$$

Proof: One starts from the definition of the pull-back and use (2.27) to get

$$(\pi^{r+1,r})^*\rho(j^{r+1}x\gamma)(\xi_1, \dots, \xi_q) = \rho(j^rx\gamma)(h\xi_1 + p\xi_1, \dots, h\xi_q + p\xi_q).$$

Then one derives the formula in the statement if one uses the definition (3.7) and the following combinatorial lemma:

Lemma 3.2 *Let L and M be linear finite dimensional spaces and $\omega : \underbrace{L \times \dots \times L}_{q\text{-times}} \rightarrow M$ a linear and antisymmetric map. Then the following formula is true:*

$$\omega(a_1 + b_1, \dots, a_q + b_q) = \varepsilon^{i_1, \dots, i_q} \sum_{k=0}^q \frac{1}{k!(q-k)!} \omega(a_{i_1}, \dots, a_{i_k}, b_{i_{k+1}}, \dots, b_{i_q}) \quad (3.10)$$

for all $a_1, \dots, a_q, b_1, \dots, b_q \in L$.

Proof: Is straightforward by induction on q . ■

One can use the proposition above to deduce that ρ is contact *iff* it verifies $h\rho = 0$ and that the expressions $p_1\rho, \dots, p_q\rho$ are contact forms in the sense of the definition given at the beginning of this Section. Finally we remind the fact that the horizontalization operation $h : \Omega^r \rightarrow \Omega^{r+1}$ verifies the properties:

$$h(\mu + \nu) = h\mu + h\nu, \quad h(\mu \wedge \nu) = h\mu \wedge h\nu \quad (3.11)$$

for any forms $\mu, \nu \in \Omega^r$,

$$hdx^i = dx^i, \quad hdy_j^\sigma = y_{j_i}^\sigma dx^i \quad (|J| \leq r) \quad (3.12)$$

and also

$$hf = (\pi^{r+1,r})^*f = f \circ \pi^{r+1,r} \quad (3.13)$$

for all smooth functions on V^r . Moreover, these three properties given above determine *uniquely* the map h .

3.2 The Structure Theorem for Contact Forms

In this subsection we give a new proof to the following fact.

Theorem 3.3 *Let (V, ψ) an adapted chart on the fibre bundle Y and let $\rho \in \Omega_q^r(Y)$, $q = 2, \dots, n$. Then ρ is contact iff it has the following expression in the associated chart:*

$$\rho = \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J + \sum_{|I|=r-1} d\omega_I^\sigma \wedge \Psi_\sigma^I \quad (3.14)$$

where $\Phi_\sigma^J \in \Omega_{q-1}^r$ and $\Psi_\sigma^I \in \Omega_{q-2}^r$ can be arbitrary forms on V^r . (We adopt the convention that $\Omega_q^r \equiv 0, \forall q < 0$).

Proof: (i) If ρ has the expression (3.14) one uses (3.11) to obtain that it is a contact form. We now concentrate on the converse statement. Firstly we need a canonical expression for any $\rho \in \Omega_q^r$. We start from the fact that the forms dx^i , ω_J^σ ($|J| \leq r-1$) and dy_I^σ ($|I|=r$) are a basis in the linear space of 1-forms. The form ρ is a polynomial of degree q in these forms (with respect to the product \wedge). We separate all the terms containing at least a factor ω_J^σ and get a decomposition

$$\rho = \rho_0 + \rho' \quad (3.15)$$

where ρ_0 has the structure

$$\rho_0 = \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J \quad (3.16)$$

and ρ' is a polynomial of degree q only in dx^i and dy_I^σ ($|I|=r$). It is clear the one can write it as follows:

$$\rho' = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s|=r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q}, \quad (3.17)$$

where $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ are smooth functions on V^r and can be assumed to verify the following symmetry property:

$$A_{\sigma_{P(1)}, \dots, \sigma_{P(s)}, i_{Q(s+1)}, \dots, i_{Q(q)}}^{I_{P(1)}, \dots, I_{P(s)}} = (-1)^{|P|+|Q|} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \quad (3.18)$$

for any permutations $P \in \mathcal{P}_s$, $Q \in \mathcal{P}_{q-s}$.

One must impose the condition $h\rho = h\rho' = 0$. Using the relations (3.11) and (3.12) one gets

$$h\rho' = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s|=r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} y_{I_1 i_1}^{\sigma_1} \dots y_{I_s i_s}^{\sigma_s} dx^{i_1} \wedge \dots \wedge dx^{i_q},$$

so the relation above is equivalent to

$$\mathcal{S}_{i_1, \dots, i_q}^- \sum_{|I_1|, \dots, |I_s|=r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} y_{I_1 i_1}^{\sigma_1} \dots y_{I_s i_s}^{\sigma_s} = 0, \quad s = 0, \dots, q. \quad (3.19)$$

Here $\mathcal{S}_{i_1, \dots, i_q}^-$ is the antisymmetrization operator in the indices i_1, \dots, i_q defined by the formula (10.8) from the Appendix.

We apply to the relation above derivative operators of the type ∂_ν^I , ($|I| = r + 1$) and obtain, taking into account the symmetry property (3.18) the following relations (see [21] , [23]):

$$A_{i_1, \dots, i_q} = 0 \quad (3.20)$$

and

$$\mathcal{S}_{i_1, \dots, i_q}^- \mathcal{S}_{I_1 p_1}^+ \cdots \mathcal{S}_{I_s p_s}^+ A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \delta_{i_1}^{p_1} \cdots \delta_{i_s}^{p_s} = 0, \quad s = 1, \dots, q. \quad (3.21)$$

(ii) We analyse in detail the relation above. This is the point where we depart from the idea of the proof from [23]. Let us introduce the following tensor spaces:

$$\mathcal{H}_s \equiv \mathcal{F}^{(-)}(\mathbb{R}^n) \otimes \underbrace{\mathcal{F}^{(+)}(\mathbb{R}^n) \otimes \cdots \otimes \mathcal{F}^{(+)}(\mathbb{R}^n)}_{s\text{-times}}$$

where $\mathcal{F}^{(\pm)}(\mathbb{R}^n)$ are the symmetric (corresp. +) and the antisymmetric (corresp. -) Fock spaces (see the Appendi). We have the well known decomposition in subspaces with fixed number of “bosons” and “fermions”:

$$\mathcal{H}_s = \bigoplus_{k=0}^r \bigoplus_{l_1, \dots, l_s \geq 0} \mathcal{H}_{k, l_1, \dots, l_s}.$$

We make the convention that $\mathcal{H}_{k, l_1, \dots, l_s} \equiv 0$ if any one of the indices k, l_1, \dots, l_s is negative or if $k > n$. Then we can consider $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ as the components of a tensor

$$A_{\sigma_1, \dots, \sigma_s} \in \mathcal{H}_{q-s, \underbrace{r, \dots, r}_{s\text{-times}}}.$$

We can write in an extremely compact way the relation (3.21) if we use the creation and the annihilation fermionic operators a^{*i}, a_i , ($i = 1, \dots, n$) and the corresponding creation and annihilation bosonic operators $b_{(\alpha)i}^*, b_{(\alpha)}^i$ ($\alpha = 1, \dots, s; i = 1, \dots, n$). We will using conventions somewhat different from that used in quantum mechanics (see the Appendix). One introduces the operators

$$B_\alpha \equiv b_{(\alpha)i}^* a^{*i}, \quad (\alpha = 1, \dots, s) \quad (3.22)$$

and proves by elementary computations that the relations (3.21) are equivalent to:

$$B_1 \cdots B_s A_{\sigma_1, \dots, \sigma_s} = 0, \quad (s = 1, \dots, q). \quad (3.23)$$

Indeed, it is elementary to check that we have for instance:

$$(B_1 A)_{\sigma_1, \dots, \sigma_s, i_0, \dots, i_l}^{\{j_0, \dots, j_k\}, I_2, \dots, I_s} = \mathcal{S}_{j_0, \dots, j_k}^+ \mathcal{S}_{i_0, \dots, i_l}^- A_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_l}^{\{j_1, \dots, j_k\}, I_2, \dots, I_s} \delta_{i_0}^{j_0}. \quad (3.24)$$

and similarly for $\alpha = 2, \dots, s$.

(iii) Equations of the type appearing in the (3.23) can be completely analysed with the help of two lemmas, which are the backbone of our paper. Firstly we analyse the case $s = q$. In this case $A_{\sigma_1, \dots, \sigma_q} \in \mathcal{H}_{0, \underbrace{r, \dots, r}_{q\text{-times}}}$.

Lemma 3.4 *Let $X \in \mathcal{H}_{0,r_1,\dots,r_q}$, $(r_1, \dots, r_q \in \mathbb{N}; \quad q = 2, \dots, n)$ verifying the relation*

$$B_1 \cdots B_q X = 0. \quad (3.25)$$

Then $X = 0$.

Proof: We use complete induction over n . For $n = 2$ we have only the case $q = 2$. The generic form of X is

$$X = \sum_{i=0}^{r_1} \sum_{j=0}^{r_2} c_{i,j} (b_{(1)1}^*)^i (b_{(1)2}^*)^{r_1-i} (b_{(2)1}^*)^j (b_{(2)2}^*)^{r_2-j} \Omega;$$

here Ω is the vacuum vector in \mathcal{H} . We extend the coefficients $c_{i,j}$ to all integer values of the indices by taking $c_{i,j} = 0$ if i (or j) is outside the set $\{1, \dots, r_1\}$ (or $\{1, \dots, r_2\}$). One computes that the relation $B_1 B_2 X = 0$ is equivalent to $c_{i-1,j} = c_{i,j-1}$; this relation gives by recurrence that $c_{i,j} = 0$, $\forall i, j \in \mathbb{Z}$ i.e. $X = 0$.

We assume the assertion from the statement to be true for a given n and we prove it for $n + 1$. In this case the indices will take the values $0, 1, \dots, n$ and we must make the substitution

$$B_\alpha \rightarrow \tilde{B}_\alpha \equiv B_\alpha + b_{(\alpha)0}^* a^{*0}, \quad (\alpha = 1, \dots, q).$$

By hypothesis we have

$$\prod_{\alpha=1}^q \tilde{B}_\alpha X = 0.$$

One immediately shows this to be equivalent to the following two equations:

$$\prod_{\alpha=1}^q B_\alpha X = 0 \quad (3.26)$$

and

$$\sum_{\alpha=1}^q (-1)^{\alpha-1} b_{(\alpha)}^{*0} B_1 \cdots \hat{B}_\alpha \cdots B_q X = 0. \quad (3.27)$$

The generic expression for X is

$$X = \sum_{t_1=0}^{r_1} \cdots \sum_{t_q=0}^{r_q} (b_{(1)}^{*0})^{t_1} \cdots (b_{(q)}^{*0})^{t_q} X_{t_1, \dots, t_q} \quad (3.28)$$

where $X_{t_1, \dots, t_q} \in \mathcal{H}_{0, r_1-t_1, \dots, r_q-t_q}$ are tensors obtained from the vacuum by applying only polynomials in the creation operators $b_{(\alpha)}^{*i}$, $(\alpha = 1, \dots, n; i = 1, \dots, n)$.

Then the equation (3.26) is equivalent to

$$\prod_{\alpha=1}^q B_\alpha X_{t_1, \dots, t_q} = 0, \forall t_1, \dots, t_q \in \mathbb{Z}. \quad (3.29)$$

If $q \leq n$ we can apply the induction hypothesis and obtain $X_{t_1, \dots, t_q} = 0, \forall t_1, \dots, t_q \in \mathbb{Z}$ i.e. $X = 0$. So it remains to study only the case $q = n + 1$. In this case one notices that the

equation (3.26) becomes an identity because $\prod_{\alpha=1}^{n+1} B_{\alpha} = 0$; indeed, in the left hand side, at least one of the operators a^{*i} , $(i = 1, \dots, n)$ appears twice. So, in this case we are left with the second equation (3.27):

$$\sum_{\alpha=1}^{n+1} (-1)^{\alpha-1} b_{(\alpha)}^{*0} B_1 \cdots \hat{B}_{\alpha} \cdots B_{n+1} X = 0. \quad (3.30)$$

One must substitute here the generic expression for X (3.28) and the following relation is produced:

$$\sum_{\alpha=1}^{n+1} (-1)^{\alpha-1} b_{(\alpha)}^{*0} B_1 \cdots \hat{B}_{\alpha} \cdots B_{n+1} X_{t_1, \dots, t_{\alpha-1}, t_{\alpha}-1, t_{\alpha+1}, \dots, t_{n+1}} = 0, \forall t_1, \dots, t_{n+1} \in \mathbf{Z}. \quad (3.31)$$

This relation can be used to prove that $X = 0$. We outline the argument. First we take $t_1 = t + 1$ ($t \geq 0$), $t_2 = \dots = t_{n+1} = 0$ in the relation above and get

$$B_2 \cdots B_{n+1} X_{\underbrace{t, 0, \dots, 0}_{n\text{-times}}} = 0.$$

The induction hypothesis can be applied and we get $X_{\underbrace{t, 0, \dots, 0}_{n\text{-times}}} = 0$, $\forall t \in \mathbf{Z}$. Similarly, one can obtain $X_{0, \dots, 0, t, 0, \dots, 0} = 0$, $\forall t \in \mathbf{Z}$ where the index t can be positioned anywhere.

Now one can prove, by induction on p that $X_{t_1, \dots, t_{n+1}} = 0$ if at least $n - p + 1$ indices are equal to zero. For $p = 1$ this statement has just been proved above. We accept it for a given p and prove it for $p + 1$. We take in (3.31) $t_{p+2} = \dots = t_{n+1} = 0$ and obtain

$$\sum_{\alpha=1}^{p+1} (-1)^{\alpha-1} b_{(\alpha)}^{*0} B_1 \cdots \hat{B}_{\alpha} \cdots B_{n+1} X_{t_1, \dots, t_{\alpha-1}, t_{\alpha}-1, t_{\alpha+1}, \dots, t_{n+1}} = 0. \quad (3.32)$$

If $t_1 = t, t_2 = t_3 = \dots = t_{p+1} = 1$ one can use the induction hypothesis to get:

$$X_{\underbrace{t, 1, \dots, 1}_{p\text{-times}}, \underbrace{0, \dots, 0}_{(n-p)\text{-times}}} = 0, \quad \forall t \in \mathbf{Z}.$$

Now one uses the relation above and (3.32) to prove that

$$X_{t_1, \dots, t_{p+1}, \underbrace{0, \dots, 0}_{(n-p)\text{-times}}} = 0; \quad (3.33)$$

this is done by recurrence on $t_2 + t_3 + \dots + t_{p+1}$.

Evidently the argument leading to the relation (3.33) works in the same way for any positioning of the $p + 1$ indices. The induction is finished and we get $X_{t_1, \dots, t_{n+1}} = 0$, $\forall t_1, \dots, t_{n+1} \in \mathbf{Z}$ i.e. $X = 0$. ∇

If we apply this lemma to the equation (3.23) for $s = q$ we get

$$A_{\sigma_1, \dots, \sigma_q}^{I_1, \dots, I_q} = 0. \quad (3.34)$$

(iv) We still have to analyse the case $0 < s < q$ of the equation (3.23). We can analyse immediately the case $s = 1$ i.e. the equation

$$B_1 X = 0 \quad (3.35)$$

if we use (10.16); we get that

$$(n + r - q + 1)X = BB^*X$$

i.e. X is of the form

$$X = BX_1 \quad (3.36)$$

for some $X_1 \in \mathcal{H}_{q-2, r-1}$.

(v) We generalize this result to all $s = 1, \dots, q - 1$.

Lemma 3.5 *Let $X \in \mathcal{H}_{k, r_1, \dots, r_s}$, $0 \leq k < n$, $0 < s \leq n$, $s + k \leq n$. Then X verifies the equation*

$$B_1 \cdots B_s X = 0 \quad (3.37)$$

iff it is of the form

$$X = \sum_{\alpha=1}^s B_\alpha X_\alpha \quad (3.38)$$

for some $X_\alpha \in \mathcal{H}_{k-1, r_1, \dots, r_{\alpha-1}, r_\alpha-1, r_{\alpha+1}, \dots, r_s}$.

Proof:

From (10.18) the implication (3.38) \Rightarrow (3.37) is obvious. We prove now the converse statement, as before, by induction on n . For $n = 2$ we can have $k = 0, 1$. In the first case the statement is true according to lemma 3.4 and the second case was analysed before at (iv). We suppose that the statement is true for a given n and we prove it for $n + 1$. With the same notations as in lemma 3.4 we have an equation of the type:

$$\prod_{\alpha=1}^s \tilde{B}_\alpha X = 0. \quad (3.39)$$

where

$$\tilde{B}_\alpha \equiv B_\alpha + b_{(\alpha)0}^* a^{*0}, \quad (\alpha = 1, \dots, q).$$

The generic form for X is

$$X = X_0 + a^{*0} Z \quad (3.40)$$

with $X_0 \in \mathcal{H}_{k, r_1, \dots, r_s}$ and $Z \in \mathcal{H}_{k-1, r_1, \dots, r_s}$ tensors obtained from the vacuum by applying only polynomials in a^{*i} ($i = 1, \dots, n$) and $b_{(\alpha)\mu}^*$ ($\mu = 0, \dots, n; \alpha = 1, \dots, s$).

The equation (3.39) becomes equivalent to the following two equations:

$$\prod_{\alpha=1}^s B_\alpha X_0 = 0 \quad (3.41)$$

and

$$\prod_{\alpha=1}^s B_{\alpha} Z = \sum_{\alpha=1}^s (-1)^{s-\alpha} b_{(\alpha)}^{*0} B_1 \cdots \hat{B}_{\alpha} \cdots B_s X_0. \quad (3.42)$$

As in lemma 3.4, the generic form of X_0 is

$$X_0 = \sum_{t_1=0}^{r_1} \cdots \sum_{t_q=0}^{r_q} \left(b_{(1)}^{*0}\right)^{t_1} \cdots \left(b_{(q)}^{*0}\right)^{t_q} X_{t_1, \dots, t_q}. \quad (3.43)$$

Then the equation (3.41) is equivalent to

$$\prod_{\alpha=1}^s B_{\alpha} X_{t_1, \dots, t_q} = 0, \quad \forall t_1, \dots, t_q \in \mathbf{Z}. \quad (3.44)$$

We have two distinct cases:

(a) $k < n$ and $s \leq n$

In this case we can apply the induction hypothesis to the relation above and obtain in the end that X_0 is of the following form:

$$X_0 = \sum_{\alpha=1}^s B_{\alpha} X_{\alpha}.$$

If we introduce this expression into the equation (3.42) we easily get:

$$\prod_{\alpha=1}^s B_{\alpha} \left(Z - \sum_{\alpha=1}^s b_{(\alpha)}^{*0} X_{\alpha} \right) = 0$$

so again we can apply the induction hypothesis to obtain that Z has the following structure:

$$Z = \sum_{\alpha=1}^s b_{(\alpha)}^{*0} X_{\alpha} + \sum_{\alpha=1}^s B_{\alpha} Z_{\alpha}.$$

Now we define

$$\tilde{X}_{\alpha} \equiv X_{\alpha} - a^{*0} Z_{\alpha}$$

and obtain from the previous relations that

$$X = \sum_{\alpha=1}^s \tilde{B}_{\alpha} \tilde{X}_{\alpha}$$

which finishes the proof.

(b) If $k = n$ then from the restrictions on s and k we necessarily have $s = 1$ and we can use (iv). If $s = n + 1$ then the same restrictions fix $k = 0$ and we can apply lemma 3.4. ∇

We can apply the lemma above to the equation (3.23) for the cases $s = 2, \dots, q - 1$ and obtain that the tensors $A_{\sigma_1, \dots, \sigma_s}$ have the following structure

$$A_{\sigma_1, \dots, \sigma_s} = \sum_{\alpha=1}^s B_{\alpha} X_{\sigma_1, \dots, \sigma_s}^{\alpha}$$

for some tensors $X_{\sigma_1, \dots, \sigma_s}^\alpha$.

Now it is the moment to use again full index notation. The relation above means that the expression $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ is a sum of terms such that every term has at least a factor of the type δ_i^j where the index j belongs to some I_p and the index i is one of the indices i_{s+1}, \dots, i_q . That's it if, say, $I_1 = \{j_1, \dots, j_r\}$ then

$$A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{\{j_1, \dots, j_r\}, \dots} = \sum_{u=1}^r \sum_{v=s+1}^q \delta_{i_v}^{j_u} \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, \hat{i}_v, \dots, i_q}^{\{j_1, \dots, \hat{j}_u, \dots, j_r\}, \dots} + \dots$$

If we substitute the preceding relation into the expression of ρ' (see (3.17)) we obtain a sum of contributions of the type

$$\tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, \hat{i}_v, \dots, i_q}^{\{j_1, \dots, \hat{j}_u, \dots, j_r\}, \dots} dy_{\{j_1, \dots, j_r\}}^{\sigma_1} \wedge dy_{I_2}^{\sigma_2} \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{j_u} \dots \wedge dx^{i_q}$$

i.e. a sum of terms containing the expression $dy_{\{j_1, \dots, j_r\}}^{\sigma_1} \wedge dx^{j_u} = -d\omega_{\{j_1, \dots, \hat{j}_u, \dots, j_r\}}^{\sigma_1}$.

So the contribution ρ' to the contact form ρ form gives the second terms from the statement of the theorem (v. the formulæ (3.15) and (3.14). ■

Let us note for further use that one can combine lemmas 3.4 and 3.5 in a single result:

Lemma 3.6 *Let $X \in \mathcal{H}_{k, r_1, \dots, r_s}$. Then X verifies the equation (3.37) iff it is of the form*

$$X = \sum_{\alpha=1}^s B_\alpha X_\alpha \tag{3.45}$$

for some $X_\alpha \in \mathcal{H}_{k-1, r_1, \dots, r_{\alpha-1}, r_\alpha-1, r_{\alpha+1}, \dots, r_s}$.

Remark 3.6.1 *One can show in fact that the decomposition of an arbitrary form given by the formulæ (3.15), (3.16) and (3.17) can be refined with the help of the so called trace decomposition identity [24]. Although we do not need this more refined decomposition we will provide an alternative proof of this fact, based on the same tricks, in the Appendix. This will emphasize once more the power of our method.*

3.3 Some Properties of the Contact Forms

We start with the transformation formula for the contact forms. We have:

Proposition 3.7 *Let (V, ψ) and $(\bar{V}, \bar{\psi})$ be two overlapping charts on Y . Then on $V^r \cap \bar{V}^r$ the following formula is true:*

$$\bar{\omega}_I^\sigma = \sum_{|J| \leq |I|} (\partial_\nu^J \bar{y}_I^\sigma) \omega_J^\nu \quad (|I| \leq r-1). \quad (3.46)$$

Proof: The proof is based on simple manipulations of the formula (2.19), the definition (3.3) of the 1-contact forms and use is also made of lemma 2.3. ■

An element $T \in \Omega_{n+1, X}^s$ is called a *differential equation* if $i_\xi T = 0$ for any $\pi^{s,1}$ -vertical vector field. In the chart V^s the differential equation T has the following expression:

$$T = T_\sigma \omega^\sigma \wedge \theta_0. \quad (3.47)$$

(see (2.35)). Using (3.46) one can indeed see that T has this form in any chart; explicitly, the transformation formula is:

$$T_\sigma = \mathcal{J}(\partial_\sigma \bar{y}^\nu) \bar{T}_\nu \quad (3.48)$$

where \mathcal{J} is the Jacobian of the chart transformation on X :

$$\mathcal{J} \equiv \det \left(\frac{\partial \bar{x}^i}{\partial x^j} \right). \quad (3.49)$$

If γ is a section of the fibre bundle $\pi : Y \rightarrow X$ then one says that *it verifies the differential equation T iff* we have

$$(j^s \gamma)^* i_Z T = 0 \quad (3.50)$$

for any vector field Z on $J^s Y$. In local coordinates we have on V^s :

$$T_\sigma \circ j^s \gamma = 0 \quad (\sigma = 1, \dots, m). \quad (3.51)$$

Another important property of the contact ideal is that it behaves naturally with respect to prolongations. More precisely, let $\pi_i : Y_i \rightarrow X_i$, $i = 1, 2$ be two fibre bundles and $\phi : Y_1 \rightarrow Y_2$ a fibre bundle morphism. Then the prolongation $j^r \phi$ (defined in the end of section 2) verifies:

$$(j^r \phi)^* \mathcal{C}(\Omega^r(Y_1)) \subset \mathcal{C}(\Omega^r(Y_2)). \quad (3.52)$$

The proof follows directly from the definition of a contact form. As a consequence, if ξ is a projectable vector field on the fibre bundle Y , then the Lie derivative of $j^r \xi$ leaves the contact ideal invariant:

$$L_{j^r \xi} \mathcal{C}(\Omega^r(Y)) \subset \mathcal{C}(\Omega^r(Y)). \quad (3.53)$$

This formula can be used to find out the explicit expression of $j^r \xi$ [2]. Indeed, if in the chart (V, ψ) we have

$$\xi = a^i(x) \frac{\partial}{\partial x^i} + b^\sigma(x, y) \partial_\sigma \quad (3.54)$$

with a^i and b^σ smooth function, then $j^r \xi$ must have the following expression in the associated chart (V^r, ψ^r) :

$$j^r \xi = a^i(x) \frac{\partial}{\partial x^i} + \sum_{|J| \leq r} b_J^\sigma \partial_\sigma^J. \quad (3.55)$$

One imposes an equivalent form of (3.53), namely

$$L_{j^r \xi} \omega_J^\sigma \in \mathcal{C}(\Omega^r(Y)), \quad |J| \leq r-1. \quad (3.56)$$

The left hand side of this relation can be computed explicitly:

$$L_{j^r \xi} \omega_J^\sigma = (d_i b_J^\sigma - b_{J_i}^\sigma - y_{Jl}^\sigma d_i a^l) dx^i + \sum_{|I|=r} (\partial_\nu^I b_J^\sigma) dy_I^\nu + \text{contact terms}.$$

The following recurrence formula for the coefficients b_J^σ follows:

$$b_{J_i}^\sigma = d_i b_J^\sigma - y_{Jl}^\sigma d_i a^l, \quad |J| \leq r-1; \quad (3.57)$$

we also have:

$$\partial_\nu^I b_J^\sigma = 0, \quad |I| = r. \quad (3.58)$$

In particular, if ξ is an evolution i.e. it has the local expression (2.30), then we have

$$j^r \xi = \sum_{|J| \leq r} (d_J \xi^\sigma) \partial_\sigma^J. \quad (3.59)$$

One may wonder what is the expression of the prolongation $j^r \phi$ (where ϕ is a bundle morphism of the fibre bundle Y). One can proceed in complete analogy with the computations above. If ϕ has the following expression in the chart (V, ψ)

$$\phi(x^i, y^\sigma) = (f^i, F^\sigma) \quad (3.60)$$

then we must have in the associated chart: (V^r, ψ^r) :

$$j^r \phi(x^i, y^\sigma, y_J^\sigma, \dots, y_{j_1, \dots, j_r}^\sigma) = (f^i, F^\sigma, F_J^\sigma, \dots, F_{j_1, \dots, j_r}^\sigma) \quad (3.61)$$

where $F_{j_1, \dots, j_k}^\sigma$, $j_1 \leq j_2 \leq \dots \leq j_k$, $k = 1, \dots, r$ are smooth functions on the chart V^r . One starts from an equivalent form of (3.52), namely:

$$(j^r \phi)^* \omega_J^\sigma \in \mathcal{C}(\Omega^r(Y)) \quad |J| \leq r-1 \quad (3.62)$$

and computes the left hand side:

$$(j^r \phi)^* \omega_J^\sigma = \left(d_l F_J^\sigma - F_{Jl}^\sigma \frac{\partial f^l}{\partial x^i} \right) dx^i + \sum_{|I|=r} (\partial_\nu^I F_J^\sigma) dy_I^\nu + \text{contact terms}, \quad |J| \leq r-1.$$

The condition (3.62) above gives us a recurrence formula for the functions F_J^σ :

$$F_{J_i}^\sigma = Q_i^l d_l F_J^\sigma \quad |J| \leq r-1; \quad (3.63)$$

we also have

$$\partial_\nu^I F_J^\sigma = 0 \quad |I| = r. \quad (3.64)$$

Let us note that the recurrence formula above formally coincide with the recurrence formula from lemma 2.3.

We close this subsection reminding another important construction appearing when one considers the so-called variationally trivial Lagrangians (they will be defined in section 7). We introduce the following subset of the space of basic forms:

$$\mathcal{J}_q^r \equiv \{\rho \in \Omega_{q,X}^r | \exists \nu \in \Omega_q^{r-1}, \quad s.t. \quad \rho = h\nu\}. \quad (3.65)$$

One notes that this subspace is closed with respect to the wedge product \wedge and also that the operator $D : \mathcal{J}_q^r \rightarrow \mathcal{J}_{q+1}^r$ given by

$$Dh\nu \equiv h d\nu \quad (3.66)$$

is well defined [3]. The operator D is called *total exterior derivative*. We list some elementary properties of this operator directly deductible from the definition.

$$D \circ D = 0 \quad (3.67)$$

and

$$Dh = hD. \quad (3.68)$$

Moreover, D is a derivation completely determined by the following relations:

$$Df = (d_i f) dx^i, \quad \forall f \in \mathcal{J}_0^r, \quad (3.69)$$

and

$$D(dx^i) = 0, \quad Dhdy_J^\sigma = 0, \quad |J| \leq r. \quad (3.70)$$

We remind the reader that we have remarked before that the operators d_i are not vector fields. However, we have the following relation, which is the next best thing except a vector field. If $f \in \Omega_0^{r-1}$ and we have two overlapping charts (V, ψ) and $(\bar{V}, \bar{\psi})$ on Y then we have on the intersection $V^r \cap \bar{V}^r$ the following relation:

$$\bar{d}_i f = Q_i^j d_j f \quad (3.71)$$

(where \bar{d}_j are the formal derivatives in the chart \bar{V}^r and the matrix Q has been defined previously: it is the inverse of the Jacobian matrix of the chart transformation - see (2.7)). The proof is elementary and consists in expressing the (globally defined) operator D in both charts.

4 Strongly Contact Forms

The concept of strongly contact form has been introduced by Krupka [21]. The idea is to observe that the definition of the contact forms is trivially satisfied if the degree of the form is $q \geq n + 1$. So, it is natural to try a generalization of the contact forms in this case. It seems plausible to use instead of the horizontalization operator h some other projection p_k from those introduced in the beginning of the preceding section. The proper definition is the following. Let $q = n + 1, \dots, N \equiv \dim(J^r Y) = m \binom{n+r}{n}$ and let $\rho \in \Omega_q^r$. One says that ρ is a *strongly contact form* iff its contact component of order $q - n$ vanishes i.e.

$$p_{q-n}\rho = 0. \quad (4.1)$$

For a certain uniformity of notations, we denote these forms by $\Omega_{q(c)}^r$. We need a structure formula for strongly contact forms, i.e. an analogue of theorem 3.3. First we need some properties of the projections p_k [21] - [23].

Lemma 4.1 *If $\rho \in \Omega_q^r$ and $\rho' \in \Omega_t^r$ then the following formula is true:*

$$p_k(\rho \wedge \rho') = \sum_{l+s=k} p_l \rho \wedge p_s \rho' \quad \forall k \geq 0 \quad (4.2)$$

where we make the convention that

$$p_k \rho \equiv 0, \quad \text{if } k > \deg(\rho), \quad \text{or } k < 0. \quad (4.3)$$

Proof: Is based on induction on q . For $q = 1$ one starts from the definition (3.7) of the operator p_k and from the definition for the wedge product, which in our case is:

$$(\rho \wedge \rho')(\xi_0, \dots, \xi_t) = \sum_{i=0}^t (-1)^i \rho(\xi_i) \rho'(\xi_0, \dots, \hat{\xi}_i, \dots, \xi_t).$$

Then, one supposes the formula true for $1, 2, \dots, q$ and proves it for $q+1$. One does not lose generality if one supposes that ρ is of the form $\rho = \rho_1 \wedge \rho_2$ with $\deg(\rho_1) = 1$ and $\deg(\rho_2) = q$ and the result for $q+1$ is obvious. ■

As a corollary, we have

Corollary 4.2 *If $\rho_i \in \Omega_{q_i}^r$, $i = 1, \dots, l$ then the following formula is true:*

$$p_k(\rho_1 \wedge \dots \wedge \rho_l) = \sum_{s_1 + \dots + s_l = k} p_{s_1} \rho_1 \wedge \dots \wedge p_{s_l} \rho_l. \quad (4.4)$$

In particular, if the order of contactness of the forms $\rho_i \in \Omega_{q_i}^r$, $i = 1, \dots, l$ is equal to 1 and if $\rho' \in \Omega_q^r$ is arbitrary, then the following formula is true:

$$p_k(\rho_1 \wedge \dots \wedge \rho_l \wedge \rho') = (\pi^{r+1,r})^* \rho_1 \wedge \dots \wedge (\pi^{r+1,r})^* \rho_l \wedge p_{k-l} \rho', \quad \forall k \geq l. \quad (4.5)$$

If $k < l$ then the right hand side is zero, according to the convention from the preceding lemma.

We still need two results from [23]. The first one is elementary.

Lemma 4.3 *Let $q \geq 1$ and $\rho \in \Omega_q^r$. Then in the associated chart (V^{r+1}, ψ^{r+1}) the following formula is valid:*

$$(\pi^{r+1,r})^* \rho = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s| \leq r} B_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \omega_{I_1}^{\sigma_1} \wedge \dots \wedge \omega_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \dots \wedge dx^{i_q} \quad (4.6)$$

where the coefficients $B_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ are smooth functions on the chart V^r and verify symmetry properties of the type (3.18). Moreover, the form $p_k \rho$ is given by the terms corresponding to $s = k$ in the sum above.

Next, we have

Lemma 4.4 *Let $q \geq 1$ and $\rho \in \Omega_q^r$. Suppose that in the associated chart (V^r, ψ^r) the form ρ has the generic expression:*

$$\rho = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s| \leq r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \dots \wedge dx^{i_q} \quad (4.7)$$

where $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ are smooth functions on V^r verifying the symmetry property (3.18). Then on the chart (V^{r+1}, ψ^{r+1}) we have

$$p_k \rho = \frac{1}{k!(q-k)!} \sum_{|I_1|, \dots, |I_k| \leq r} B_{\sigma_1, \dots, \sigma_k, i_{k+1}, \dots, i_q}^{I_1, \dots, I_k} \omega_{I_1}^{\sigma_1} \wedge \dots \wedge \omega_{I_k}^{\sigma_k} \wedge dx^{i_{k+1}} \dots \wedge dx^{i_q} \quad (4.8)$$

where

$$B_{\sigma_1, \dots, \sigma_k, i_{k+1}, \dots, i_q}^{I_1, \dots, I_k} = \mathcal{S}_{i_{k+1}, \dots, i_q}^- \sum_{s=k}^q \binom{q-k}{q-s} \sum_{|I_{k+1}|, \dots, |I_s| \leq r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} y_{I_{k+1} i_{k+1}}^{\sigma_{k+1}} \dots y_{I_s i_s}^{\sigma_s}. \quad (4.9)$$

Proof: We use the definition (3.4) to write

$$\begin{aligned} (\pi^{r+1,r})^* \rho &= \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s| \leq r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \\ &(\omega_{I_1}^{\sigma_1} + y_{I_1 i_1}^{\sigma_1} dx^{i_1}) \wedge \dots \wedge (\omega_{I_s}^{\sigma_s} + y_{I_s i_s}^{\sigma_s} dx^{i_s}) \wedge dx^{i_{s+1}} \dots \wedge dx^{i_q} \end{aligned} \quad (4.10)$$

and now we can apply lemma 3.2 with $L \mapsto \Omega^r$, $M \mapsto \Omega_s^r$ and $\omega \mapsto \Lambda$ where

$$\Lambda(\omega_{I_1}^{\sigma_1}, \dots, \omega_{I_s}^{\sigma_s}) = \sum_{|I_1|, \dots, |I_s| \leq r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \omega_{I_1}^{\sigma_1} \dots \wedge \omega_{I_s}^{\sigma_s}.$$

Then simple rearrangements leads to the formula from the statement. \blacksquare

Now we can give the structure theorem for strongly contact forms. As in the preceding section, the proof will be based on Fock space machinery and will differ from the original proof from [21].

Theorem 4.5 Let $n+1 \leq q \leq N$ and $\rho \in \Omega_q^r$. Let (V, ψ) be a chart on Y . Then ρ is a strongly contact form iff it has the following expression in the associated chart (V^r, ψ^r) :

$$\rho = \sum_{p+s=q-n+1} \sum_{|J_1|, \dots, |J_p| \leq r-1} \sum_{|I_1|, \dots, |I_s| = r-1} \omega_{J_1}^{\sigma_1} \cdots \wedge \omega_{J_p}^{\sigma_p} \wedge d\omega_{I_1}^{\nu_1} \cdots \wedge d\omega_{I_s}^{\nu_s} \wedge \Phi_{\sigma_1, \dots, \sigma_p, \nu_1, \dots, \nu_s}^{J_1, \dots, J_p, I_1, \dots, I_s} \quad (4.11)$$

where $\Phi_{\sigma_1, \dots, \sigma_p, \nu_1, \dots, \nu_s}^{J_1, \dots, J_p, I_1, \dots, I_s}$ are differential forms of degree $n-1-s$ on V^r . (This imposes that the first sum runs in fact only for $s \leq n-1$).

Proof: If ρ has the expression from the statement, the corollary above gives us $p_{q-n}\rho = 0$. We prove the converse statement by induction on q .

(i) Let $q = n+1$ and $\rho \in \Omega_{n+1}^r$ such that $p_1\rho = 0$. We start from the same decomposition of the form ρ as in theorem 3.3 i.e. that given by the formulæ (3.15)-(3.17). Using (3.15) and the corollary above we have from the preceding equation:

$$\sum_{|J| \leq r-1} \omega_J^\sigma \wedge h\Phi_\sigma^J + p_1\rho' = 0. \quad (4.12)$$

But the preceding lemma gives us the following very explicit formula for the second contribution:

$$p_1\rho' = \frac{1}{(q-1)!} \sum_{|I|=r} B_{\sigma, i_2, \dots, i_q}^I \omega_I^\sigma \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_q}$$

where

$$B_{\sigma_1, i_2, \dots, i_q}^{I_1} = \mathcal{S}_{i_2, \dots, i_q}^- \sum_{s=1}^q \binom{n}{n+1-q} \sum_{|I_2|, \dots, |I_s| = r-1} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} y_{I_2 i_2}^{\sigma_2} \cdots y_{I_s i_s}^{\sigma_s}.$$

Because the 1-form ω_I^σ appears only in the second term of (4.12) the two terms must vanish separately i.e. we have

$$p_1\rho' = 0. \quad (4.13)$$

and

$$\sum_{|J| \leq r-1} \omega_J^\sigma \wedge h\Phi_\sigma^J = 0. \quad (4.14)$$

From (4.13) we get

$$B_{\sigma, i_2, \dots, i_q}^I = 0$$

which can be transformed, as in the proof of theorem 3.3 into

$$B_2 \cdots B_s A_{\sigma_1, \dots, \sigma_s} = 0 \quad (s = 1, \dots, q).$$

In fact, because of the symmetry property (3.18) we have:

$$B_1 \cdots \hat{B}_\alpha \cdots B_s A_{\sigma_1, \dots, \sigma_s} = 0 \quad (\alpha = 1, \dots, s; s = 1, \dots, q). \quad (4.15)$$

A consequence of this relation is

$$B_1 \cdots B_s A_{\sigma_1, \dots, \sigma_s} = 0 \quad (s = 1, \dots, q)$$

which implies (see lemmas 3.4 and 3.5)) that

$$A_{\sigma_1, \dots, \sigma_q} = 0 \quad (4.16)$$

and

$$A_{\sigma_1, \dots, \sigma_s} = \sum_{\alpha=1}^s B_\alpha A_{\sigma_1, \dots, \sigma_s}^\alpha \quad (s = 1, \dots, q-1) \quad (4.17)$$

for some tensors $A_{\sigma_1, \dots, \sigma_s}^\alpha$.

If we substitute the expression of $A_{\sigma_1, \dots, \sigma_s}$ above into the initial equation (4.15) we get immediately

$$B_1 \cdots B_s A_{\sigma_1, \dots, \sigma_s}^\alpha = 0$$

so lemma 3.5 can again be applied to produce the following expression

$$A_{\sigma_1, \dots, \sigma_s} = \sum_{\alpha, \beta=1}^s B_\alpha B_\beta A_{\sigma_1, \dots, \sigma_s}^{\alpha\beta} \quad (s = 1, \dots, q-1) \quad (4.18)$$

for some tensors $A_{\sigma_1, \dots, \sigma_s}^{\alpha\beta} = 0$. The last expression identically verifies the equation (4.15) so it is the general solution of it. As in the end of theorem 3.3 it is the time to revert to full index notations. Because in the formula above we have *two* B -type operators we will obtain that the functions $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ are sums of terms containing *two* delta factors, so in the end *two* factors of the type $d\omega_I^\nu$ we show up. Explicitly, ρ' must necessarily have the following structure:

$$\rho' = \sum_{|I_1|=|I_2|=r-1} d\omega_{I_1}^{\nu_1} \wedge d\omega_{I_2}^{\nu_2} \wedge \Phi_{\nu_1 \nu_2}^{I_1 I_2}. \quad (4.19)$$

On the other hand, it is easy to see that (4.14) is equivalent to

$$h\Phi_\sigma^J = 0$$

and theorem 3.3 can be applied. Combining with the formula above, we obtain the structure formula from the statement (4.11) for $q = n+1$.

(ii) We suppose that if $p_{q-n}\rho = 0$ then ρ' has the expression (4.11) for $q' = n+1, \dots, q-1$ and we prove the same statement for q . So we have

$$p_{q-n}\rho = 0.$$

Because ρ is a polynomial of degree q (with respect to the wedge product \wedge) in the differentials ω_J^σ $|J| \leq r-1$, dy_I^σ , $|I| = r$ and dx^i one can write it uniquely as follows:

$$\rho = \sum_{s=0}^q \frac{1}{q!} \sum_{|J_1|, \dots, |J_s| \leq r-1} \omega_{J_1}^{\sigma_1} \cdots \wedge \omega_{J_s}^{\sigma_s} \wedge \Phi_{\sigma_1, \dots, \sigma_s}^{J_1, \dots, J_s} \quad (4.20)$$

where $\Phi_{\sigma_1, \dots, \sigma_s}^{J_1, \dots, J_s}$ are polynomials of degree $q-s$ in the differentials dy_I^σ , $|I| = r$ and dx^i . Using the corollary above one obtains the following equation:

$$\sum_{s=0}^q \frac{1}{q!} \sum_{|J_1|, \dots, |J_s| \leq r-1} \omega_{J_1}^{\sigma_1} \cdots \wedge \omega_{J_s}^{\sigma_s} \wedge p_{q-n-s} \Phi_{\sigma_1, \dots, \sigma_s}^{J_1, \dots, J_s} = 0$$

which is equivalent to:

$$p_{q-n-s}\Phi_{\sigma_1,\dots,\sigma_s}^{J_1,\dots,J_s} = 0 \quad (s = 0, \dots, q). \quad (4.21)$$

For $s = 1, \dots, q$ one can apply the induction hypothesis and obtain that the forms $\Phi_{\sigma_1,\dots,\sigma_s}^{J_1,\dots,J_s}$ are sums of the type (4.11). It remains to analyse the case $s = 0$ i.e. the equation

$$p_{q-n}\Phi = 0 \quad (4.22)$$

with Φ having a structure similar to (3.17):

$$\Phi = \sum_{s=q-n}^q \frac{1}{s!(q-s)!} \sum_{|I_1|,\dots,|I_s|=r} A_{\sigma_1,\dots,\sigma_s,i_{s+1},\dots,i_q}^{I_1,\dots,I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q}. \quad (4.23)$$

Using the preceding lemma one obtains that

$$p_{q-n}\Phi = \frac{1}{(q-n)!n!} \sum_{|I_1|,\dots,|I_{q-n}|=r} B_{\sigma_1,\dots,\sigma_{q-n},i_{q-n+1},\dots,i_q}^{I_1,\dots,I_{q-n}} \omega_{I_1}^{\sigma_1} \wedge \dots \wedge \omega_{I_{q-n}}^{\sigma_{q-n}} \wedge dx^{i_{q-n+1}} \dots \wedge dx^{i_q}$$

where

$$B_{\sigma_1,\dots,\sigma_{q-n},i_{q-n+1},\dots,i_q}^{I_1,\dots,I_{q-n}} = \mathcal{S}_{i_{q-n+1},\dots,i_q}^- \sum_{s=q-n}^q \binom{n}{q-s} \sum_{|I_{q-n+1}|,\dots,|I_s|=r} A_{\sigma_1,\dots,\sigma_s,i_{s+1},\dots,i_q}^{I_1,\dots,I_s} y_{I_{q-n+1}}^{\sigma_{q-n+1}} \dots y_{I_s}^{\sigma_s}. \quad (4.24)$$

The condition on Φ translates into

$$B_{\sigma_1,\dots,\sigma_{q-n},i_{q-n+1},\dots,i_q}^{I_1,\dots,I_{q-n}} = 0$$

and this can be shown to be equivalent to

$$B_{q-n+1} \dots B_s A_{\sigma_1,\dots,\sigma_s} = 0 \quad (s = q-n, \dots, q).$$

In fact, because of the symmetry property (3.18) we have more generally:

$$B_{\alpha_1} \dots B_{\alpha_{s-q+n}} A_{\sigma_1,\dots,\sigma_s} = 0 \quad \forall \alpha_1, \dots, \alpha_{s-q+n}, (s = q-n, \dots, q). \quad (4.25)$$

This relation can be investigated following the ideas from (i) (see rel. (4.15)) and the general solution of (4.25) can be found in the form:

$$A_{\sigma_1,\dots,\sigma_s} = \sum_{\alpha_1,\dots,\alpha_{q-n+1}=1}^s B_{\alpha_1} \dots B_{\alpha_{q-n+1}} A_{\sigma_1,\dots,\sigma_s}^{\alpha_1,\dots,\alpha_{q-n+1}} \quad (4.26)$$

for some tensors $A_{\sigma_1,\dots,\sigma_s}^{\alpha_1,\dots,\alpha_{q-n+1}}$.

If we use full index notations, this time $q-n+1$ factors of the type $d\omega_I^\rho$, $|I| = r-1$ will appear in every term of Φ . Collecting all terms we get for ρ the formula (4.11). \blacksquare

5 Euler-Lagrange and Helmholtz-Sonin forms

An interesting problem in differential geometry is the following one. Suppose we have a differential form ρ on a given manifold Y . What other (globally defined) differential forms can be attached to it? This problem can be rigorously formulated [19] and the answer is that there is essentially only one possibility, namely the exterior differential $d\rho$ of ρ . In other words the condition of correct behaviour with respect to all possible charts transformations limits drastically the possible solution to this kind of problem. But what happens when the manifold Y has a supplementary structure, say is a fibre bundle? Then, there will be some restrictions on the charts transformation so other solutions can appear. In this section we will prove that in this case indeed new possibilities can appear, as for instance the Euler-Lagrange and Helmholtz-Sonin form. We will follow essentially [2] making the observation that much of the line of the argument can be adapted from infinite jet bundle extensions to our case i.e. finite bundle extensions.

5.1 Lie-Euler Operators

The central combinatorial trick used in [2] to prove the existence of the Euler-Lagrange form is the concept of *total differential operator* which, by definition, is any linear map $P : \mathcal{E}(J^r Y) \rightarrow \Omega^s$ covering the identity map $id : J^r Y \rightarrow J^r Y$ with $s \geq r$. (One considers, of course, $\mathcal{E}(J^r Y)$ and $J^r Y$ as fibre bundles over $J^r Y$). We will consider in the following that s is sufficiently great; in fact one needs that $s > 2r + 2$. Suppose that ξ is an evolution having the local structure (2.30) in the chart (V^r, ψ^r) . Then the image $P(\xi) \in \Omega^s$ must have the expression:

$$P(\xi) = \sum_{|I| \leq r} (d_I \xi^\sigma) P_\sigma^I = \sum_{k=0}^r (d_{j_1} \cdots d_{j_k} \xi^\sigma) P_I^{j_1, \dots, j_k}, \quad (5.1)$$

where P_σ^I are (local) differential forms in the chart (V^s, ψ^s) and, as usual, $d_j = d_j^s$ (see (2.20)).

Then one has the following combinatorial lemma [2]:

Lemma 5.1 *In the conditions above, the following formula is true:*

$$P(\xi) = \sum_{|I| \leq r} d_I (\xi^\sigma Q_\sigma^I) \quad (5.2)$$

where

$$Q_\sigma^I \equiv \sum_{|J| \leq r - |I|} (-1)^{|J|} \binom{|I| + |J|}{|I|} d_J P_\sigma^{IJ} \quad (5.3)$$

and one assumes that the action of a formal derivative d_j on a form is realized by its action on the function coefficients.

Proof: One starts from the right hand side of (5.2) and uses Leibnitz rule:

$$\sum_{|I| \leq r} d_I (\xi^\sigma Q_\sigma^I) = \sum_{|I| \leq r} \sum_{(J,K)} (d_J \xi^\sigma) (d_K Q_\sigma^I)$$

where the sum over (J, K) is over all partitions of the set I . One can rearrange this as follows:

$$\begin{aligned} \sum_{|I| \leq r} d_I(\xi^\sigma Q_\sigma^I) &= \sum_{|J|+|K| \leq r} \binom{|J|+|K|}{|J|} (d_J \xi^\sigma) (d_K Q_\sigma^{JK}) = \\ &= \sum_{|J| \leq r} (d_J \xi^\sigma) \sum_{|K| \leq r-|J|} \binom{|J|+|K|}{|J|} d_K Q_\sigma^{JK}. \end{aligned}$$

Now one proves by elementary computations that

$$P_\sigma^I = \sum_{|J| \leq r-|I|} \binom{|J|+|K|}{|J|} d_J \xi^\sigma Q_\sigma^{IJ}. \quad (5.4)$$

and that finishes the proof. \blacksquare

Remark 5.1.1 *One notices that the relation (5.2) uniquely determines the forms Q_σ^I .*

We proceed now to formulate the main result of this subsection. The proof is an easy adaptation to the finite jet bundle extension case of the proof from [2].

Theorem 5.2 *Let $q \geq n$ and $P : \mathcal{E}(J^r Y) \rightarrow \Omega_{q,X}^s$ a total differential operator. Let (V, ψ) and $(\bar{V}, \bar{\psi})$ two overlapping charts on Y and let us construct on the intersection of the corresponding associated charts the forms Q_σ^I and \bar{Q}_σ^I according to the preceding lemma. Then the following relation is true on the intersection $V^s \cap \bar{V}^s$:*

$$Q_\sigma = (\partial_\sigma \bar{y}^\nu) \bar{Q}_\nu. \quad (5.5)$$

In particular, there exists a globally defined form, denoted by $E(P)(\xi)$ such that in the chart (V^r, ψ^r) we have

$$E(P)(\xi) = Q_\sigma \xi^\sigma. \quad (5.6)$$

Proof: The generic expression for Q_σ^I in the chart V^s is (see (2.35)):

$$Q_\sigma^I = \sum_{|J_1|, \dots, |J_l| \leq s} Q_{\sigma, \nu_1, \dots, \nu_l}^{I, J_1, \dots, J_l} dy_{J_1}^{\nu_1} \wedge \dots \wedge dy_{J_l}^{\nu_l} \wedge \theta_0. \quad (5.7)$$

Here $l = q - n$ and $Q_{\sigma, \nu_1, \dots, \nu_l}^{I, J_1, \dots, J_l}$ are smooth functions on V^s having appropriate antisymmetry properties.

In the other chart \bar{V}^s we have a similar expression:

$$\begin{aligned} \bar{Q}_\sigma^I &= \sum_{|J_1|, \dots, |J_l| \leq s} \bar{Q}_{\sigma, \nu_1, \dots, \nu_l}^{I, J_1, \dots, J_l} d\bar{y}_{J_1}^{\nu_1} \wedge \dots \wedge d\bar{y}_{J_l}^{\nu_l} \wedge \bar{\theta}_0 = \\ &= \mathcal{J} \sum_{|J_1|, \dots, |J_l| \leq s} \tilde{Q}_{\sigma, \nu_1, \dots, \nu_l}^{I, J_1, \dots, J_l} dy_{J_1}^{\nu_1} \wedge \dots \wedge dy_{J_l}^{\nu_l} \wedge \theta_0 \end{aligned} \quad (5.8)$$

where \mathcal{J} is the Jacobian of the chart transformation on X (see (3.49)).

If we define for any I with $|I| \leq r-1$

$$R_\sigma^I \equiv i_{d_j} Q_\sigma^{jI}$$

then, using the formula above, it is easy to prove that:

$$Q_\sigma^{j_1, \dots, j_k} = \mathcal{S}_{j_1, \dots, j_k}^+ dx^{j_1} \wedge R_\sigma^{j_2, \dots, j_k} = \frac{1}{k} \sum_{p=1}^k dx^{j_p} \wedge R_\sigma^{j_1, \dots, \hat{j}_p, \dots, j_k}. \quad k = 0, \dots, r.$$

As a consequence one can rewrite the local formula (5.2) as follows:

$$P(\xi) = \xi^\sigma Q_\sigma + dx^i \wedge d_i R(\xi), \quad (5.9)$$

where we have defined

$$R(\xi) \equiv \sum_{|I| \leq r-1} d_I(\xi^\sigma R_\sigma^I). \quad (5.10)$$

So, in the overlap $V^s \cap \bar{V}^s$ we have

$$\xi^\sigma Q_\sigma - \bar{\xi}^\sigma \bar{Q}_\sigma = d\bar{x}^i \wedge \bar{d}_i \bar{R}(\bar{\xi}) - dx^i \wedge d_i R(\xi).$$

Let us remark now that from the definition of the forms $Q_\sigma^{j_1, \dots, j_k}$ (see (5.3)) it follows that its function coefficients depend only on the variables $(x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_{2r-k}}^\sigma)$; as a consequence, the function coefficients of the form $R(\xi)$ depend only on the variables $(x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_{2r-1}}^\sigma)$. In this case we can apply formula (3.71) to the preceding relation and we obtain:

$$\xi^\sigma Q_\sigma - \bar{\xi}^\sigma \bar{Q}_\sigma = dx^i \wedge d_i \tilde{R}(\xi)$$

where

$$\tilde{R}(\xi) = \bar{R}(\bar{\xi}) - R(\xi) = \sum_{|J_1|, \dots, |J_l| \leq s} \tilde{R}_{\nu_1, \dots, \nu_l}^{i, J_1, \dots, J_l}(\xi) dy_{J_1}^{\nu_1} \wedge \dots \wedge dy_{J_l}^{\nu_l} \wedge \theta_i;$$

here we have defined

$$\theta_i \equiv (-1)^{i-1} x^1 \wedge \dots \wedge dx^{i-1} \wedge dx^{i+1} \wedge \dots \wedge dx^n \quad (5.11)$$

and $\tilde{R}_{\nu_1, \dots, \nu_l}^{i, J_1, \dots, J_l}(\xi)$ are smooth functions on the overlap $V^s \cap \bar{V}^s$.

If we also use (2.30) we obtain

$$\xi^\sigma \left[Q_{\sigma, \nu_1, \dots, \nu_l}^{\emptyset, J_1, \dots, J_l} - \mathcal{J}(\partial_\sigma \bar{y}^\zeta) \bar{\xi}^\sigma \tilde{Q}_{\zeta, \nu_1, \dots, \nu_l}^{\emptyset, J_1, \dots, J_l} \right] = d_i \tilde{R}_{\nu_1, \dots, \nu_l}^{i, J_1, \dots, J_l}(\xi).$$

Now one proves that both sides are zero in a standard way: one picks a section γ with support in $W \subset \pi(V) \cap \pi(\bar{V})$ such that the closure \bar{W} of W is compact, takes ξ^σ with support in the open set $U \subset \bar{W}$ and integrates on \bar{W} the following relation (which follows from the preceding one):

$$\xi^\sigma \left[Q_{\sigma, \nu_1, \dots, \nu_l}^{\emptyset, J_1, \dots, J_l} - \mathcal{J}(\partial_\sigma \bar{y}^\zeta) \bar{\xi}^\sigma \tilde{Q}_{\zeta, \nu_1, \dots, \nu_l}^{\emptyset, J_1, \dots, J_l} \right] \circ j^s \gamma = d_i \tilde{R}_{\nu_1, \dots, \nu_l}^{i, J_1, \dots, J_l}(\xi) \circ j^s \gamma.$$

Use of Stokes theorem is made and of the arbitrariness of γ and it follows that:

$$Q_{\sigma, \nu_1, \dots, \nu_l}^{\emptyset, J_1, \dots, J_l} = \mathcal{J}(\partial_\sigma \bar{y}^\zeta) \bar{\xi}^\sigma \tilde{Q}_{\zeta, \nu_1, \dots, \nu_l}^{\emptyset, J_1, \dots, J_l}$$

If we introduce this equality in (5.7) and (5.8) we obtain the relation (5.5). ■

The operator $E(P)$ defined by (5.6) is called the *Euler operator* associated to the total differential operator P ; it has the local expression:

$$E(P)(\xi) = \xi^\sigma E_\sigma(P) \quad (5.12)$$

where

$$E_\sigma(P) = \sum_{|I|=0}^r (-1)^{|I|} d_I P_\sigma^I. \quad (5.13)$$

Now one takes $\lambda \in \Omega_{n,X}^r$ and constructs the total differential operator P_λ :

$$P_\lambda(\xi) \equiv L_{pr(\xi)} \lambda. \quad (5.14)$$

Suppose that λ has the local expression (2.32). Then lemma 5.1 can be applied and gives the following formula:

$$P_\lambda(\xi) = \sum_{|I|=0}^r d_I \left(\xi^\sigma E_\sigma^I(L) \right) \theta_0 \quad (5.15)$$

where

$$E_\sigma^I(L) \equiv \sum_{|J| \leq r-|I|} (-1)^{|J|} \binom{|I|+|J|}{|I|} d_J \partial_\sigma^{IJ} L \quad (5.16)$$

are the so-called *Lie-Euler operators*.

In particular,

$$Q_\sigma = E_\sigma(L) \theta_0 \quad (5.17)$$

and the Euler operator associated to P_λ has the following expression:

$$E(P_\lambda) = \xi^\sigma E_\sigma(L) \theta_0 \quad (5.18)$$

where

$$E_\sigma(L) \equiv \sum_{|J| \leq r} (-1)^{|J|} d_J \partial_\sigma^J L \quad (5.19)$$

are the *Euler-Lagrange expressions*.

The theorem above leads to

Proposition 5.3 *If $\lambda \in \Omega_{n,X}^r$ is a Lagrange form, then there exists a globally defined $n+1$ -form, denoted by $E(\lambda)$ such that we have in the chart V^s :*

$$E(\lambda) = E_\sigma(L) \omega^\sigma \wedge \theta_0. \quad (5.20)$$

Proof: Indeed, one combines the expression (5.17) with the transformation properties (3.46) and (5.5) to obtain

$$E_\sigma(L) = \mathcal{J}(\partial_\sigma \bar{y}^\nu) \bar{E}_\nu(\bar{L}); \quad (5.21)$$

the globallity of $E(\lambda)$ follows immediately. ■

One calls this form the *Euler-Lagrange form* associated to λ and notes that it is a differential equation (see the beginning of subsection 3.3 more precisely the formula (3.47)). In general, a differential equation $T \in \Omega_{n+1,X}^s$ is called *(locally) variational* or *of Euler-Lagrange type* iff there exists a (local) Lagrange form $\lambda \in \Omega_{n,X}^r$ ($s \geq 2r$) such that

$$T = E(\lambda). \quad (5.22)$$

One notices that in this case the general form of a differential equation (3.51) coincides with the well-known form of the Euler-Lagrange equations.

Remark 5.3.1 *There are other ways of proving the globallity of the Euler-Lagrange form. One can use the existence of the so-called Lepagean equivalents [18], but the combinatorial analysis seems to be more complicated. Also, an argument based on the connection between the action functional and the Euler-Lagrange expression is available [21]*

5.2 Some Properties of the Euler-Lagrange Form

We collect for further use some properties of the Euler-Lagrange form. We follow, essentially [2], with the appropriate modifications.

First, one finds rather easily from (5.21) that the Euler-Lagrange form behaves naturally with respect to bundle morphisms. More precisely, if $\phi \in \text{Diff}(Y)$ is a bundle morphism, then one has:

$$(j^s \phi)^* E(\lambda) = E((j^r \phi)^* \lambda) \quad (5.23)$$

for any Lagrange form λ . From here, we obtain by differentiation:

$$L_{j^s \xi} E(\lambda) = E(L_{j^r \xi} \lambda) \quad (5.24)$$

for any projectable vector field ξ on Y .

Now we have

Lemma 5.4 *If f is a smooth function on V^r then we have in V^s , $s > 2r + 2$ the following formulæ:*

$$E_\sigma^{Ij}(d_I f) = \mathcal{S}_{Ij}^+ \delta_I^j E_\sigma^I(f), \quad |I| = 0, \dots, r \quad (5.25)$$

and

$$E_\sigma(d_I f) = 0. \quad (5.26)$$

Proof: From lemma 5.1 we have in V^s :

$$\sum_{|I| \leq r} (d_I \xi^\sigma) (\partial_\sigma^I L) = \sum_{|I| \leq r} d_I (\xi^\sigma E_\sigma^I(L))$$

for any smooth function L on V^r . We make $r \rightarrow r+1$ and $L \rightarrow d_j f$ and we have:

$$\sum_{|I| \leq r+1} (d_I \xi^\sigma) \left(\partial_\sigma^I d_j f \right) = \sum_{|I| \leq r+1} d_I \left(\xi^\sigma E_\sigma^I (d_j f) \right).$$

The left hand side can be rewritten using the commutation formula (2.23) and one obtains:

$$\sum_{k=1}^{r+1} d_{i_1} \dots d_{i_k} \left(\xi^\sigma \mathcal{S}_{i_1, \dots, i_k}^+ \delta_j^{i_1} E_\sigma^{i_2, \dots, i_k} (f) \right) = \sum_{k=0}^{r+1} d_{i_1} \dots d_{i_k} \left(\xi^\sigma E_\sigma^{i_1, \dots, i_k} (d_j f) \right).$$

Using remark 5.1.1 we obtain the relations from the statement. ■

Corollary 5.5 *Let A^I , $|I| = l \geq 1$ be some smooth functions on V^r and $f \equiv d_I A^I$. Then we have on V^s , $s > 2(r+l)$:*

$$E_\sigma^J(f) = 0, \quad |J| \leq |I| - 1. \quad (5.27)$$

Corollary 5.6 *Let ξ be an evolution on Y and $\lambda \in \Omega_{n,X}^r$ a Lagrange form. Then the following formula is true:*

$$E(L_{j^r \xi} \lambda) = E(i_{j^s \xi} E(\lambda)). \quad (5.28)$$

Proof: One has by direct computation and use on lemma 5.1:

$$E(L_{j^r \xi} \lambda - i_{j^s \xi} E(\lambda)) = \sum_{|I|=1}^r E_\nu \left(d_I (\xi^\sigma E_\sigma^I(L)) \right) \omega^\nu \wedge \theta_0;$$

but the right hand side is zero, according to the preceding corollary. ■

Corollary 5.7 *Let $\lambda \in \Omega_{n,X}^r$ a Lagrange form. Then the following formula is true:*

$$L_{j^s \xi} E(\lambda) = E(i_{j^s \xi} E(\lambda)). \quad (5.29)$$

Indeed, one combines the preceding corollary with (5.24) and obtains this formula.

5.3 Helmholtz-Sonin Forms

In this section, we follow an idea of [5] to prove the existence of the (globally) defined Helmholtz-Sonin form. We have the following central result.

Theorem 5.8 *Let $T \in \Omega_{n+1,X}^s$ be a differential equation with the local form given by (3.47). We define the following expressions in any chart V^t , $t > 2s$:*

$$H_{\sigma\nu}^J \equiv \partial_\nu^J T_\sigma - (-1)^{|J|} E_\sigma^J(T_\nu), \quad |J| \leq s. \quad (5.30)$$

Then there exists a globally defined $(n+2)$ -form, denoted by $H(T)$ such that in any chart V^t we have:

$$H(T) = \sum_{|J| \leq s} H_{\sigma\nu}^J \omega_J^\nu \wedge \omega^\sigma \wedge \theta_0. \quad (5.31)$$

Proof: (i) We begin with a construction from [5]. Let ξ be an evolution; we define a (global) n -form $H_\xi(T)$ according to:

$$H_\xi(T) \equiv L_{j^s \xi} T - E(i_{j^s \xi} T) \quad (5.32)$$

Elementary computations and use of corollary 5.5 leads to the following local expression:

$$H_\xi(T) = \sum_{|I| \leq s} (d_I \xi^\sigma) H_{\sigma\nu}^I \omega^\sigma \wedge \theta_0. \quad (5.33)$$

(ii) Now one determines the transformation formula at a change of charts for the expressions $d_I \xi^\sigma$. One considers the evolution ξ on Y and writes the expression of the vector field $pr(\xi)$ on the overlap $V^t \cap \bar{V}^t$; the following formula easily emerges:

$$\bar{d}_I \bar{\xi}^\sigma = \sum_{|J| \leq |I|} \left(\partial_\nu^J \bar{y}_I^\sigma \right) (d_J \xi^\nu), \quad |I| = 0, \dots, s. \quad (5.34)$$

(iii) Using the transformation formula (5.34) one can obtain the transformation formula for the expressions $H_{\sigma\nu}^I$: one has in the overlap $V^t \cap \bar{V}^t$:

$$H_{\mu\zeta}^J = \mathcal{J} \sum_{|I| \geq |J|} \left(\partial_\mu^J \bar{y}_I^\nu \right) (\partial_\zeta \bar{y}^\sigma) \bar{H}_{\sigma\nu}^I. \quad (5.35)$$

This transformation formula should now be combined with (3.46) and one obtains that $H(T)$ has an invariant meaning; we have

$$\sum_{|I| \leq s} \bar{H}_{\sigma\nu}^I \bar{\omega}_I^\nu \wedge \bar{\omega}^\sigma \wedge \bar{\theta}_0 = \sum_{|J| \leq s} H_{\mu\zeta}^J \omega_J^\mu \wedge \omega^\zeta \wedge \theta_0$$

on $V^t \cap \bar{V}^t$ and the proof is finished. ■

$H(T)$ is called the *Helmholtz-Sonin form* associated to T and $H_{\sigma\nu}^I$ are the *Helmholtz-Sonin expressions* associated to T .

A well-known corollary of the theorem above is:

Corollary 5.9 *The differential equation T is locally variational iff $H(T) = 0$.*

Proof: (i) If there exists a Lagrange form λ such that locally $T = E(\lambda)$, then from (5.29) it follows that $H_\xi(T) = 0$ for any evolution ξ . This implies that the Helmholtz-Sonin expressions associated to T are zero i.e. $H(E(\lambda)) = 0$.

(ii) The converse of this statement is done by some explicit construction. Suppose that T is such that $H(T) = 0$. Then we define the following Lagrangian on the chart V^s :

$$L = \int_0^1 dt y^\sigma T_\sigma \circ \chi_t^s, \quad \chi_t^s(x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_s}^\sigma) = (x^i, ty^\sigma, ty_j^\sigma, \dots, ty_{j_1, \dots, j_s}^\sigma). \quad (5.36)$$

Then one proves by direct computation that $E_\sigma(L) = \mathcal{T}_\sigma$ so, T is locally variational. ■

The (local) expression (5.36) is called the *Tonti-Vainberg* Lagrangian.

6 The Exact Variational Sequence

This section is divided in two parts. The first one includes in a brief way the standard proof of the exactness of the variational sequence following the lines of [21]. The second part is devoted to the characterization of some elements of the variational sequence by (globally) defined forms. As it is pointed out in [21], this can be done using the Euler-Lagrange and the Helmholtz-Sonin forms defined previously. In this part more details are given because the proofs from the literature are rather sketchy. In particular, the proof we offer for the characterization of the $n + 1$ term in the variational sequence by the Helmholtz-Sonin form is new.

6.1 The Exactness of the Variational Sequence

If $\pi : Y \rightarrow X$ is a fibre bundle and $U, V \subset Y$ are charts such that $U \subset V$, we denote by $\iota_{U,V} : U^r \rightarrow V^r$ the canonical inclusion. Then, the collection $\{\Omega_q^r(V)\}$ ($q \geq 0$), $\{\iota_{U,V}^*\}$ is a presheaf denoted by Ω_q^r . Next, one introduces the subspaces $\theta_q^r \in \Omega_{q(c)}^r$ by:

$$\theta_1^r \equiv \Omega_{1(c)}^r, \quad \theta_q^r \equiv d\Omega_{q-1(c)}^r + \Omega_{q(c)}^r \quad (q = 2, \dots, N = \dim(J^r Y)); \quad (6.1)$$

this is also a presheaf and one can easily verify that:

$$\theta_q^r = \Omega_{q(c)}^r \quad (q = 2, \dots, n), \quad \theta_q^r = 0 \quad (q > P), \quad d\theta_q^r \subset \theta_{q+1}^r; \quad P \equiv m \binom{n+r-1}{n} + 2(n-1). \quad (6.2)$$

Next, one introduces the so-called contact homotopy operator. The construction is the following. Let $U \subset \mathbb{R}^n$ (resp. $V \subset \mathbb{R}^m$) an open set (resp. a ball centred in $0 \in \mathbb{R}^m$) and $W = U \times V$. One considers the operator χ_r as a map $\chi_t^r : [0, 1] \times J^r W \rightarrow J^r W$ given by:

$$\chi_r(t, (x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_s}^\sigma)) = (x^i, ty^\sigma, ty_j^\sigma, \dots, ty_{j_1, \dots, j_s}^\sigma). \quad (6.3)$$

Then for any $\rho \in \Omega_q^r(W)$ we have the *unique* decomposition

$$(\chi_t^r)^* \rho = dt \wedge \rho_0(t) + \rho_1(t) \quad (6.4)$$

where $\rho_0(t)$ (resp. $\rho_1(t)$) are $q - 1$ (resp. q) forms which do not contain the differential dt . Then the *contact homotopy operator* is by definition the map $A : \Omega_q^r(V) \rightarrow \Omega_{q-1}^r(V)$ given by:

$$A\rho \equiv \int_0^1 \rho_0(t). \quad (6.5)$$

Moreover one has:

$$\rho_1(1) = \rho, \quad \rho_1(0) = (\tau_r)^*(\zeta_0)^* \rho \quad (6.6)$$

where $\tau_r : V^r \rightarrow V$ is the canonical projection on the first component:

$$\tau_r(x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_s}^\sigma) \equiv (x^i)$$

and $\zeta_0 : U \rightarrow J^0 W$ is the zero section given by:

$$\zeta_0(x^i) \equiv (x^i, 0, \dots, 0).$$

Then we have (see [21]):

Lemma 6.1 (i) Let $\rho \in \Omega^r W$ be arbitrary. The following formula is true:

$$\rho = Ad\rho + dA\rho + (\tau_r)^*(\zeta_0)^*\rho \quad (6.7)$$

(ii) If ρ is contact, then:

$$\rho = Ad\rho + dA\rho \quad (6.8)$$

and

$$p_{k-1}A\rho = Ap_k\rho \quad (k = 1, \dots, q). \quad (6.9)$$

Proof: (i) The proof of the first formula is standard. First one finds out from (6.4) that:

$$(\chi_r)^*d\rho = d_0\rho_1 + dt \wedge \left(\frac{\partial \rho_1}{\partial t} - d_0\rho_0 \right)$$

where d_0 is the exterior differentiation with respect to all variables except t . This produces by integration:

$$Ad\rho = \rho_1(1) - \rho_1(0) - dA\rho$$

and using (6.6) we find the formula.

(ii) For the first relation one uses the structure theorem 3.3 for the contact forms and sees that last term in (6.7) is zero. For the second relation one applies $(id \times \pi^{r+1,r})^*$ to the relation (6.4) and after some simple manipulations one gets:

$$(\pi^{r+1,r})^*\rho_0 = \sum_{k=0}^q (p_k\rho)_0 \Rightarrow (\pi^{r+1,r})^*A\rho = \sum_{k=0}^q p_k A\rho = \sum_{k=1}^q Ap_k\rho.$$

Now use is made of lemma 4.4 and the second formula follows. \blacksquare

The central result of [21] follows. We will insist only on the part of the proof which can be simplified with the Fock space tricks.

Theorem 6.2 We consider the maps $d: \theta_q^r \longrightarrow \theta_{q+1}^r$; then the long sequence of sheaves

$$0 \longrightarrow \theta_1^r \xrightarrow{d} \theta_2^r \xrightarrow{d} \dots \xrightarrow{d} \theta_P^r \longrightarrow 0 \quad (6.10)$$

is exact.

Proof:

(i) The exactness in θ_1^r is elementary [21]. If $\beta \in \theta_1^r = \Omega_{1(c)}^r$ then the structure theorem 3.3 can be used to write in V^r :

$$\beta = \sum_{|J| \leq r-1} \Phi_\sigma^J \omega_J^\sigma$$

with Φ_σ^J some smooth functions on V^r . Then

$$d\beta = - \sum_{|J|=r-1} \Phi_\sigma^J dy_{J_i}^\sigma \wedge dx^i + \dots$$

where by \dots we mean terms without the differentials dy_I^σ ($|I| = r$); so, the closedness of β gives us $\Phi_\sigma^J = 0$ ($|J| = r - 1$). This means that, in fact, in the expression above of β the sum finishes at $r - 2$. Continuing by recurrence we arrive at $\beta = 0$.

(ii) We prove the exactness in θ_q^r , ($q = 2, \dots, q$). If $\beta \in \theta_q^r = \Omega_{q(c)}^r$ then again we can apply the structure theorem 3.3 to write (after simple rearrangements):

$$\beta = \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \tilde{\Phi}_\sigma^J + \sum_{|I|=r-1} d\omega_I^\sigma \wedge \Psi_\sigma^I = \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J + \sum_{|I|=r-1} d(\omega_I^\sigma \wedge \Psi_\sigma^I)$$

so the closedness condition is

$$\sum_{|J| \leq r-1} d\omega_J^\sigma \wedge \Phi_\sigma^J - \sum_{|J| \leq r-1} \omega_J^\sigma \wedge d\Phi_\sigma^J = 0.$$

Applying p_1 to this relation we find out with lemma 4.1:

$$\sum_{|J| \leq r-1} d\omega_J^\sigma \wedge h\Phi_\sigma^J - \sum_{|J| \leq r-1} \omega_J^\sigma \wedge hd\Phi_\sigma^J = 0. \quad (6.11)$$

One uses now for Φ_σ^J the standard form

$$\Phi_\sigma^J = \chi_\sigma^J + \zeta_\sigma^J$$

where χ_σ^J is generated by ω_K^σ , ($|K| \leq r - 1$) and ζ_σ^J is a polynomial of degree $q - 1$ in dx^i and dy_I^σ , ($|I| = r - 1$):

$$\zeta_{\nu_1}^{I_1} = \sum_{s=1}^q \frac{1}{s!(q-s)!} \sum_{|I_2|, \dots, |I_s|=r} A_{\nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} dy_{I_2}^{\nu_2} \wedge \dots \wedge dy_{I_s}^{\nu_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q}$$

where the symmetry properties of the type (3.18) leave out the indices I_1 and ν_1 . Then (6.11) becomes:

$$\sum_{|J|=r-1} dy_{J_i}^\sigma \wedge dx^i \wedge h\zeta_\sigma^J + \dots = 0$$

where by \dots we mean an expression which does not contain the differentials dy_I^σ , ($|I| = r$). This relations leads to

$$\mathcal{S}_{I_i}^+ dx^i \wedge h\zeta_\sigma^I = 0;$$

like in the proof of theorem 3.3 this is equivalent to

$$\mathcal{S}_{I_1 i_1}^+ \dots \mathcal{S}_{I_s i_s}^+ \mathcal{S}_{i_1, \dots, i_q}^- \delta_{i_1}^{p_1} \dots \delta_{i_s}^{p_s} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} = 0 \quad (s = 1, \dots, q),$$

or, in tensor notations:

$$B_1 \dots B_s A_{\nu_1, \dots, \nu_s} = 0 \quad (s = 1, \dots, q).$$

Using the usual argument, it follows that the forms ζ_σ^I ($|I| = r - 1$) are generated by the differentials $d\omega_K^\sigma$, ($|K| = r - 1$). It follows that Φ_σ^J , ($|J| = r - 1$) is are contact forms. By

recurrence it follows that Φ_σ^J , ($|J| \leq r-1$) is are contact forms. This information must be inserted back in the initial expression of the form β ; one has that

$$\beta = \beta_0 + d\gamma$$

where β_0 (resp. γ) are 2-contact (resp. contact) forms. The closedness condition reduces now to $d\beta_0 = 0$ and (6.8) can be applied; one gets $\beta_0 = dA\beta_0$ so finally it follows that β is given by $\beta = d(A\beta_0 + \gamma)$. But, using (6.9) one gets that $h(A\beta_0 + \gamma) = Ap_1\beta_0 = 0$ i.e. $A\beta_0 + \gamma \in \theta_{q-1}^r$. In other words $\beta \in \text{Im}(d)$.

(iii) The proof of the exactness in θ_q^r ($q > n$) is also standard [21]. Let $\beta \in \theta_q^r$ i.e. $\beta = \beta_0 + d\gamma$ where $\beta_0 \in \Omega_{q(c)}^r$ and $\gamma \in \Omega_{q-1(c)}^r$ such that $d\beta = 0$. Then $d\beta_0 = 0$ and we can apply (6.8) to obtain $\beta_0 = dA\beta_0$. As a consequence $\beta = d(A\beta_0 + \gamma)$. But formula (6.9) also implies: $p_{q-1-n}(A\beta_0 + \gamma) = Ap_{q-n}\beta_0 = 0$ so in fact $A\beta_0 + \gamma \in \Omega_{q-1(c)}^r \subset \theta_{q-1}^r$; this gives $\beta \in d\theta_{q-1}^r \subset \text{Im}(d)$. ■

This theorem has the following consequence.

Theorem 6.3 *Let $E_q : \Omega_q^r / \theta_q^r \rightarrow \Omega_{q+1}^r / \theta_{q+1}^r$ be given by*

$$E_q([\rho]) \equiv [d\rho] \quad (6.12)$$

where $[\rho]$ is the class of ρ modulo θ_q^r . Then the quotient sequence

$$0 \longrightarrow \mathbb{R} \longrightarrow \Omega_0^r \xrightarrow{E_0} \Omega_1^r / \theta_1^r \xrightarrow{E_1} \dots \xrightarrow{E_{P-1}} \Omega_P^r / \theta_P^r \xrightarrow{E_P} \Omega_{P+1}^r \xrightarrow{d} \dots \xrightarrow{d} \Omega_N^r \longrightarrow 0 \quad (6.13)$$

is a acyclic resolution of the constant sheaf \mathbb{R} . In particular it is exact.

One calls this the *variational sequence of order r over Y* and denotes for simplicity: $\mathcal{V}_q^r \equiv \Omega_q^r / \theta_q^r$.

Some special classes have distinct names. So, if $\lambda \in \Omega_n^r$ then the class $[\lambda] \in \Omega_{n+1}^r / \theta_{n+1}^r$ is called the *Euler-Lagrange class* of λ . If $T \in \Omega_{n+1}^r$ then $[T] \in \Omega_{n+2}^r / \theta_{n+2}^r$ is called the *Helmholtz-Sonin class* of T .

Let us note in the end of this subsection that for any $q = n+1, \dots, P$, $s > r$ there exists a canonical isomorphism

$$i_{s,r} : \Omega_q^r / \theta_q^r \rightarrow \text{Im} \left(\tau_q^s \circ (\pi^{s,r+1})^* \circ p_{q-n} \right),$$

where $\tau_q^s : \Omega_q^s \rightarrow \Omega_q^s / \theta_q^s$ is the canonical projection. The explicit expression is

$$i_{s,r}([\rho]) = \tau_q^s \circ (\pi^{s,r+1})^* \circ p_{q-n}(\rho) \quad (6.14)$$

and this definition is consistent [21].

6.2 Characterisation of the Variational Sequence by Forms

In this subsection we give characterizations of \mathcal{V}_q^r , $q = n, n+1$ using Euler-Lagrange and Helmholtz-Sonin forms. First we have

Theorem 6.4 *If $\lambda \in \Omega_{n,X}^r$ is any Lagrange form, then we have for any $s > 2r$*

$$i_{s,r}(E_n([\lambda])) = [E(\lambda)] \quad (6.15)$$

where $E(\lambda)$ is the Euler-Lagrange form associated to λ (see (5.19) and (5.20)).

Proof: We take the proof from [21]. If in the chart (V^r, ψ^r) the expression of λ is given by (2.32), then we have on (V^s, ψ^s) :

$$(\pi^{s,r})^* d\lambda = \sum_{k=0}^r \left(\partial_\sigma^{j_1, \dots, j_k} L \right) \omega_{j_1, \dots, j_k}^\sigma \wedge \theta_0.$$

Now we note that using (5.11) we have

$$d(\omega_{j_1, \dots, j_{k-1}}^\sigma \wedge \theta_{j_k}) = -\omega_{j_1, \dots, j_k}^\sigma \wedge \theta_0$$

for $k \geq 1$ so the preceding relation can be rearranged as follows:

$$(\pi^{s,r})^* d\lambda = (\partial_\sigma L) \omega^\sigma \wedge \theta_0 + \sum_{k=1}^r \left(d_{j_k} \partial_\sigma^{j_1, \dots, j_k} L \right) \omega_{j_1, \dots, j_{k-1}}^\sigma \wedge \theta_0 + dF_1 + G_1$$

where F_1 (resp. G_1) is a contact (resp. 2-contact form).

One can iterate the procedure and proves by induction on l that:

$$\begin{aligned} (\pi^{s,r})^* d\lambda &= \left[\sum_{k=0}^l (-1)^k \left(d_{j_1} \cdots d_{j_k} \partial_\sigma^{j_1, \dots, j_k} L \right) \right] \omega^\sigma \wedge \theta_0 + \\ &+ (-1)^l \sum_{k=l+1}^r \left(d_{j_{k-l+1}} \cdots d_{j_k} \partial_\sigma^{j_1, \dots, j_k} L \right) \omega_{j_1, \dots, j_{k-l}}^\sigma \wedge \theta_0 + dF_l + G_l \quad (l = 1, 2, \dots, r) \end{aligned}$$

where F_l (resp. G_l) is a contact (resp. 2-contact form).

In particular if we take $l = r$ we get

$$(\pi^{s,r})^* d\lambda = \left[\sum_{k=0}^r (-1)^k \left(d_{j_1} \cdots d_{j_k} \partial_\sigma^{j_1, \dots, j_k} L \right) \right] \omega^\sigma \wedge \theta_0 + dF_r + G_r = E_\sigma(L) \omega^\sigma \wedge \theta_0 + dF_r + G_r$$

i.e.

$$(\pi^{s,r})^* d\lambda = E_\sigma(L) \omega^\sigma \wedge \theta_0 \pmod{\theta_{n+1}^r}.$$

This is exactly the formula from the statement. \blacksquare

Similarly we have:

Theorem 6.5 *Let $T \in \Omega_{n+1,X}^s$ be a differential equation. The for any $t > 2s$ the following formula is true:*

$$i_{t,s}(E_{n+1}([T])) = \left[\frac{1}{2} H(T) \right] \quad (6.16)$$

where $H(T)$ is the Helmholtz-Sonin form associated to T .

Proof:

Suppose that in the chart (V^s, ψ^s) we have the local expression (3.47) for T ; then we have on (V^t, ψ^t) :

$$(\pi^{t,s})^* dT = \sum_{k=0}^s \left(\partial_\nu^{j_1, \dots, j_k} T_\sigma \right) \omega_{j_1, \dots, j_k}^\nu \wedge \omega^\sigma \wedge \theta_0.$$

Using the same trick as in the preceding theorem we can rewrite this as follows:

$$\begin{aligned} (\pi^{t,s})^* dT &= \partial_\nu T_\sigma \omega^\sigma \wedge \omega^\nu \wedge \theta_0 - \sum_{k=1}^s \left(d_{j_k} \partial_\nu^{j_1, \dots, j_k} T_\sigma \right) \omega^\sigma \wedge \omega_{j_1, \dots, j_{k-1}}^\nu \wedge \theta_0 \\ &\quad - \sum_{k=1}^s \left(\partial_\nu^{j_1, \dots, j_k} T_\sigma \right) \omega_{j_k}^\sigma \wedge \omega_{j_1, \dots, j_{k-1}}^\nu \wedge \theta_0 + dF_1 + G_1 \end{aligned}$$

where F_1 (resp. G_1) is 2-contact (resp. 3-contact).

We want to use the same idea as in the preceding theorem. We have to find the proper induction hypothesis. After some thought this proves to be:

$$\begin{aligned} (\pi^{t,s})^* dT &= \sum_{k=0}^l (-1)^p \sum_{p=k}^l \binom{p}{k} \left(d_{j_{k+1}} \cdots d_{j_p} \partial_\nu^{j_1, \dots, j_p} T_\sigma \right) \omega_{j_1, \dots, j_k}^\sigma \wedge \omega^\nu \wedge \theta_0 + \\ &+ (-1)^l \sum_{k=l+1}^s \sum_{p=k-l}^k \binom{l}{k-p} \left(d_{j_{p+1}} \cdots d_{j_k} \partial_\nu^{j_1, \dots, j_k} T_\sigma \right) \omega_{j_{k-l+1}, \dots, j_p}^\sigma \wedge \omega_{j_1, \dots, j_{k-l}}^\nu \wedge \theta_0 + dF_l + G_l \end{aligned}$$

where F_l (resp. G_l) is 2-contact (resp. 3-contact) and $l = 1, \dots, s$.

The induction from l to $l+1$ is accomplished with the same trick only the computations become much more involved. For $l = s$ the preceding formula gives:

$$(\pi^{t,s})^* dT = \sum_{k=0}^s (-1)^k \sum_{p=k}^l (-1)^{p-k} \binom{p}{k} \left(d_{j_{k+1}} \cdots d_{j_p} \partial_\nu^{j_1, \dots, j_p} T_\sigma \right) \omega_{j_1, \dots, j_k}^\sigma \wedge \omega^\nu \wedge \theta_0 + dF_s + G_s$$

where F_s (resp. G_s) is 2-contact (resp. 3-contact). Using the definition of the Lie-Euler expressions (5.16) we can write this formula in a more compact way:

$$(\pi^{t,s})^* dT = \sum_{|J| \leq s} (-1)^{|J|} E_\sigma^J(T_\sigma) \omega_J^\sigma \wedge \omega^\nu \wedge \theta_0 + dF_s + G_s.$$

We add this result to the initial expression for $(\pi^{t,s})^* dT$ and divide by 2. If we use the definition for the Helmholtz-Sonin forms (5.30) we obtain:

$$(\pi^{t,s})^* dT = \frac{1}{2} H(T) + dF_s + G_s = \frac{1}{2} H(T) \pmod{\theta_{n+2}^r}. \quad \blacksquare$$

7 Variationally Trivial Lagrangians

A *variationally trivial Lagrange form of order r* is any $\lambda \in \Omega_{n,X}^r$ such that $E(\lambda) = 0$. In other words, the corresponding Euler-Lagrange expressions are identically zero, or the Euler-Lagrange equations are identities for any section γ . Here we give the general form of such a Lagrange form. We follow, essentially [22] and [3] but as before we simplify considerably the proofs using the techniques developed in Section 3.2.

First, we remind the reader that $\rho \in \Omega_q^{r+1}$ is called $\pi^{r+1,r}$ -projectable iff there exists $\rho' \in \Omega_q^r$ such that

$$\rho = (\pi^{r+1,r})^* \rho'. \quad (7.1)$$

One can easily see that locally this amounts to the condition that if ρ is written as a polynomial in dx^i and dy_J^σ ($|J| \leq r+1$) then, the differentials dy_J^σ ($|J| = r+1$) must be absent and moreover, the coefficient functions must not depend on y_J^σ ($|J| = r+1$).

We start with the following result [21]:

Proposition 7.1 *Let $\eta \in \Omega_q^r$ ($q = 1, \dots, n-1$) such that $hd\eta$ is a $\pi^{r+1,r}$ -projectable $(q+1)$ -form. Then one can write η as follows:*

$$\eta = \nu + d\phi + \psi \quad (7.2)$$

where $\nu \in \Omega_{q,X}^r$ is a basic q -form and $\psi \in \Omega_{q(c)}^r$ is a contact q -form.

Proof:

Let (V, ψ) be a chart on Y . Then in the associated chart (V^r, ψ^r) one can write η in the standard form

$$\eta = \eta_0 + \eta_1 \quad (7.3)$$

where in η_0 we collect all terms containing at least one factor ω_J^σ ($|J| \leq r-1$) and η_1 is a polynomial only in dx^i and dy_J^σ ($|J| = r+1$):

$$\eta_1 = \sum_{s=0}^q \sum_{|I_1|, \dots, |I_s|=r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q}, \quad (7.4)$$

where the coefficients $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ have antisymmetry properties of type (3.18).

In particular η_0 is a contact form so we have $hd\eta = hd\eta_1$ i.e. the form $hd\eta_1$ is, by hypothesis, $\pi^{r+1,r}$ -projectable. One first computes in general:

$$\begin{aligned} d\eta_1 = & \sum_{s=0}^q \sum_{|I_1|, \dots, |I_s|=r} \sum_{|J| \leq r-1} \left(\partial_\nu^J A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \right) \omega_J^\nu \wedge dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q} \\ & + \sum_{s=0}^{q+1} \sum_{|I_1|, \dots, |I_s|=r} \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_{q+1}}, \end{aligned}$$

where

$$\begin{aligned} \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, I_s} &\equiv \frac{1}{s} \sum_{k=1}^s (-1)^{k-1} \partial_{\sigma_k}^{I_k} A_{\sigma_1, \dots, \hat{\sigma}_k, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, \hat{I}_k, \dots, I_s} + \\ &\frac{1}{q+1-s} \sum_{k=s+1}^{q+1} (-1)^{k-1} d_{i_k} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, \hat{i}_k, \dots, i_{q+1}}^{I_1, \dots, I_s}. \end{aligned}$$

or, equivalently

$$\begin{aligned} \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, I_s} &\equiv \mathcal{S}_{(I_1, \sigma_1), \dots, (I_s, \sigma_s)}^- \partial_{\sigma_1}^{I_1} A_{\sigma_2, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_2, \dots, I_s} + \\ &(-1)^s \mathcal{S}_{i_{s+1}, \dots, i_{q+1}}^- d_{i_{s+1}} A_{\sigma_1, \dots, \sigma_s, i_{s+2}, \dots, i_{q+1}}^{I_1, \dots, I_s} \end{aligned} \quad (7.5)$$

where $\mathcal{S}_{(I_1, \sigma_1), \dots, (I_s, \sigma_s)}^-$ is the antisymmetrization projector in the corresponding couples of indices. Then one gets

$$hd\eta = h d\eta_1 = \sum_{s=0}^{q+1} \sum_{|I_1|, \dots, |I_s|=r} \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, I_s} y_{I_1 i_1}^{\sigma_1} \cdots y_{I_s i_s}^{\sigma_s} dx^{i_1} \wedge \cdots \wedge dx^{i_{q+1}}$$

which is $\pi^{r+1, r}$ -projectable *iff* the coefficient functions do not depend on y_I^σ ($|I| = r+1$).

(ii) By repeatedly applying derivative operators, one obtains, as in [12], that the condition of $\pi^{r+1, r}$ -projectability is equivalent to

$$\mathcal{S}_{i_1, \dots, i_{q+1}}^- \mathcal{S}_{I_1 p_1}^+ \cdots \mathcal{S}_{I_s p_s}^+ \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, I_s} \delta_{i_1}^{p_1} \cdots \delta_{i_s}^{p_s} = 0, \quad (s = 1, \dots, q+1)$$

or, using a familiar argument from [12]:

$$B_1 \cdots B_s \tilde{A}_{\sigma_1, \dots, \sigma_s} = 0, \quad (s = 1, \dots, q+1). \quad (7.6)$$

It is possible to apply lemma 3.6 and one obtains the following generic expression:

$$\tilde{A}_{\sigma_1, \dots, \sigma_s} = \sum_{\alpha=1}^s B_\alpha \tilde{A}_{\sigma_1, \dots, \sigma_s}^\alpha$$

or, in index notations, this means the function $\tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_1, \dots, I_s}$ is a sum of terms containing at least a factor of the type δ_i^j where j belongs to one of the multi-indices I_p , $p = 1, \dots, s$ and $i \in \{i_{s+1}, \dots, i_{q+1}\}$.

If we insert back into the expression of $d\eta_1$ we get after minor prelucrations that

$$d\eta_1 = \nu_1 + d\phi + \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J \quad (7.7)$$

with $\nu_1 \in \Omega_{q, X}^r$ a basic q -form, $\phi \in \Omega_{q(c)}^r$ and Φ_σ^J a polynomial of degree q in dx^i and dy_I^σ $|I| = r$.

This relation implies that

$$d\nu_1 + \sum_{|J| \leq r-1} (d\omega_J^\sigma \wedge \Phi_\sigma^J - \omega_J^\sigma \wedge d\Phi_\sigma^J) = 0. \quad (7.8)$$

One applies the operator p_1 to this equality and uses (4.1); a new relation is obtained, namely:

$$p_1 d\nu_1 + \sum_{|J| \leq r-1} \left(d\omega_J^\sigma \wedge h\Phi_\sigma^J - \omega_J^\sigma \wedge h d\Phi_\sigma^J \right) = 0. \quad (7.9)$$

One must consider now the generic forms for ν_1 and Φ_σ^J namely:

$$\nu_1 = A_{i_1, \dots, i_{q+1}} dx^{i_1} \wedge \dots \wedge dx^{i_{q+1}} \quad (7.10)$$

and

$$\Phi_\sigma^J = \sum_{s=0}^q \sum_{|I_1|, \dots, |I_s|=r} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{J, I_1, \dots, I_s} dy_{I_1}^{\nu_1} \wedge \dots \wedge dy_{I_s}^{\nu_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q}; \quad (7.11)$$

here we can assume a (partial) symmetry property of the type (3.18):

$$\Phi_{\sigma, \nu_{P(1)}, \dots, \nu_{P(s)}, i_{Q(s+1)}, \dots, i_{Q(q)}}^{J, I_{P(1)}, \dots, I_{P(s)}} = (-1)^{|P|+|Q|} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{J, I_1, \dots, I_s} \quad (7.12)$$

and we make the convention that

$$\Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{J, I_1, \dots, I_s} = 0, \quad \forall J \text{ s.t. } |J| \geq r. \quad (7.13)$$

These expression must be plugged into the equation (7.9). One finds out from this equation the following consequence:

$$\mathcal{S}_{i_1, \dots, i_{q+1}}^- \mathcal{S}_{I_1 P_1}^+ \dots \mathcal{S}_{I_s P_s}^+ \left[\tilde{\Phi}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_{q+1}}^{\{j_1, \dots, j_k\}, I_1, \dots, I_s} + (-1)^s \mathcal{S}_{j_1, \dots, j_k} \delta_{i_{s+1}}^{j_1} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+2}, \dots, i_{q+1}}^{\{j_2, \dots, j_k\}, I_1, \dots, I_s} \right] \times \\ \delta_{i_1}^{p_1} \dots \delta_{i_s}^{p_s} = 0, \quad (s = 1, \dots, q+1, k = 0, \dots, r). \quad (7.14)$$

Here we have defined, in analogy with (7.5):

$$\tilde{A}_{\sigma_0, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_0, \dots, I_s} \equiv \mathcal{S}_{(I_1, \sigma_1), \dots, (I_s \sigma_s)}^- \partial_{\sigma_1}^{I_1} A_{\sigma_0, \sigma_2, \dots, \sigma_s, i_{s+1}, \dots, i_{q+1}}^{I_0, I_2, \dots, I_s} + \\ (-1)^s \mathcal{S}_{i_{s+1}, \dots, i_{q+1}}^- d_{i_{s+1}} A_{\sigma_0, \dots, \sigma_s, i_{s+2}, \dots, i_{q+1}}^{I_0, I_1, \dots, I_s}. \quad (7.15)$$

To simplify the analysis of the relation (7.14), one observes that it is possible to define a map

$$\Delta' : \oplus_{s=0}^q \mathcal{H}_{q-s, k, \underbrace{r, \dots, r}_{s\text{-times}}} \rightarrow \oplus_{s=0}^{q+1} \mathcal{H}_{q+1-s, k, \underbrace{r, \dots, r}_{s\text{-times}}}$$

according to

$$(\Delta' \Phi)_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_{q+1}}^{J, I_1, \dots, I_s} = \tilde{\Phi}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_{q+1}}^{J, I_1, \dots, I_s}. \quad (7.16)$$

Then the equation above takes the form:

$$B_1 \dots B_s \left((\Delta \Phi)_s^k + B_0 \Phi_s^{k-1} \right) = 0, \quad (s = 1, \dots, q+1, k = 0, \dots, r) \quad (7.17)$$

where $\Phi_s^k \in \mathcal{H}_{q-s, k, \underbrace{r, \dots, r}_{s\text{-times}}}$ is the tensor of components $\Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{J, I_1, \dots, I_s}$, $|J| = k$.

We remind the reader that we have made the convention

$$\Phi_s^{-1} = 0, \quad \Phi_s^r = 0. \quad (7.18)$$

To solve the preceding equation one first establishes by direct computation that

$$\{\Delta', B_0\} = 0. \quad (7.19)$$

and

$$(\Delta')^2 = 0. \quad (7.20)$$

Now we can solve the system (7.17) by a procedure similar to the descent procedure applied in the BRST quantization scheme.

We take in (7.17) $k = r$ and, taking into account the convention (7.18), we obtain:

$$B_0 \cdots B_s \Phi_s^{r-1} = 0, \quad (s = 1, \dots, q). \quad (7.21)$$

Lemma 3.6 can be applied and it follows that we have a generic expression of the following form:

$$\Phi_s^{r-1} = \sum_{\alpha=0}^s B_\alpha \Phi_s^{r-1, \alpha}, \quad (s = 1, \dots, q).$$

If we substitute this expression into the initial expression (7.11) of Φ_σ^J , $|J| = r - 1$ we find out that the contributions corresponding to $\alpha = 1, \dots, s$ are producing contact terms. So, it appears that we can redefine the expressions Φ_σ^J such that instead of (7.7) we have

$$d\eta_1 = \nu_1 + d\phi + \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J + \psi \quad (7.22)$$

with ψ some 2-contact form and moreover

$$\Phi_s^{r-1} = B_0 C_s^{r-1}, \quad (s = 1, \dots, q) \quad (7.23)$$

for some tensors $C_s^{r-1} \in \mathcal{H}_{q-s-1, r-2, \underbrace{r, \dots, r}_{s\text{-times}}}$.

The procedure can now be iterated taking into (7.17) $k = r - 1$, etc. The result of this descent procedure is the following: one can redefine the tensors Φ_σ^J such that we have (7.22) and

$$\Phi_s^k = B_0 C_s^k + (\Delta C^{k+1})_s, \quad (s = 1, \dots, q; \quad k = 0, \dots, r - 1) \quad (7.24)$$

for some tensors $C_s^k \in \mathcal{H}_{q-s-1, k-1, \underbrace{r, \dots, r}_{s\text{-times}}}$ by convention

$$C_s^0 = 0, \quad C_s^r = 0, \quad (s = 1, \dots, q).$$

In full index notations this means that we have

$$\begin{aligned} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{\{j_1, \dots, j_k\}, I_1, \dots, I_s} &= (-1)^s \mathcal{S}_{j_1, \dots, j_k}^+ \mathcal{S}_{i_{s+1}, \dots, i_q}^- \delta_{i_{s+1}}^{j_1} C_{\sigma, \nu_1, \dots, \nu_s, i_{s+2}, \dots, i_q}^{\{j_2, \dots, j_k\}, I_1, \dots, I_s} + \\ &\quad \tilde{C}'_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_q}^{\{j_1, \dots, j_k\}, I_1, \dots, I_s} \quad (s = 1, \dots, q; k = 0, \dots, r). \end{aligned} \quad (7.25)$$

Now one can substitute this expression for the tensors Φ_{\dots} into the original expression for the forms (7.11). After some algebra one obtains that

$$\sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J = \sum_{|J| \leq r-1} \Phi_{\sigma, i_1, \dots, i_q}^J \omega_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_q} + d\phi_1 \quad (7.26)$$

where $\phi_1 \in \Omega_{q(c)}^r$ is a contact form.

It emerges that the relation (7.22) becomes:

$$d\eta_1 = \nu_1 + d\phi_1 + \sum_{|J| \leq r-1} \Phi_{\sigma, i_1, \dots, i_q}^J \omega_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_q} + \psi \quad (7.27)$$

where ϕ_1 (resp. ψ) is some contact (resp. 2-contact) form of degree q (resp. $q+1$).

(iii) The last step of our proof consists in using the contact homotopy operator A (see the definition contained in the relations (6.3) - (6.5)). We apply the relation (6.7) to the form $\rho = \eta_1 - \phi_1$:

$$\eta_1 - \phi_1 = A \left(\nu_1 + \sum_{|J| \leq r-1} \Phi_{\sigma, i_1, \dots, i_q}^J \omega_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_q} + \psi \right) + dA(\eta_1 - \phi_1) + \eta_1(0) - \phi_1(0). \quad (7.28)$$

One proves by direct computation the following facts:

- $A\nu_1 = 0$;
- if ψ is 2-contact form, then $A\psi$ is a contact form;
- $A(\sum \dots)$ and $\eta_1(0)$ are basic forms;
- if ϕ_1 is a contact form, then $\phi_1(0)$ is also a contact form.

Inserting this information in the preceding relation one obtains that the result from the statement of the proposition is true for the form η_1 . Taking into account (7.3) we obtain the same result for the form η . ■

Let us note that the converse of this statement is not true. In fact the condition of $\pi^{r+1, r}$ -projectability imposes additional constraints on the basic form ν which will be analysed in the next lemma. A complete proof of the following result appears in [3]; here we offer an alternative proof which seems to be much more simpler.

Proposition 7.2 *Let $\nu \in \Omega_{q, X}^r$ ($q = 1, \dots, n-1$) Then the basic form $h\nu$ is $\pi^{r+1, r}$ -projectable iff there exists for any chart (V, ψ) a form $\tilde{\nu}_V \in \Omega_q^{r-1}$ such that we have in the associated chart (V^r, ψ^r) the equality $\nu = h\tilde{\nu}_V$.*

Proof: According to (2.31) we have in the associated chart (V^r, ψ^r) :

$$\nu = A_{i_1, \dots, i_q} dx^{i_1} \wedge \dots \wedge dx^{i_q}.$$

Then one obtains by direct computation that:

$$h\nu = \left[\left(d_{i_0} A_{i_1, \dots, i_q} \right) + \sum_{|I|=r} \left(\partial_\sigma^I A_{i_1, \dots, i_q} \right) y_{I i_0}^\sigma \right] dx^{i_0} \wedge \dots \wedge dx^{i_q}.$$

(here d_j is the formal derivative on V^r).

This form is $\pi^{r+1,r}$ -projectable *iff* the square bracket does not depend on y_I^σ ($|I| = r+1$), i.e.

$$\mathcal{S}_{i_0, \dots, i_q}^- \mathcal{S}_{Ip}^+ \delta_{i_0}^p \left(\partial_\sigma^I A_{i_1, \dots, i_q} \right) = 0. \quad (7.29)$$

It is clear that this relation is very similar to those already analysed. To be able to apply the central result contained in lemma 3.6 one must do a little trick. From the previous relation we obtain by derivation:

$$\mathcal{S}_{i_0, \dots, i_q}^- \mathcal{S}_{Ip0}^+ \delta_{i_0}^{p0} \left(\partial_{\sigma_0}^{I_0} \dots \partial_{\sigma_q}^{I_q} A_{i_1, \dots, i_q} \right) = 0. \quad (7.30)$$

If we define the tensor $A_{\sigma_0, \dots, \sigma_q} \in \mathcal{H}_q, \underbrace{r, \dots, r}_{(q+1)\text{-times}}$ by

$$A_{\sigma_0, \dots, \sigma_q, i_1, \dots, i_q}^{I_0, \dots, I_q} \equiv \partial_{\sigma_0}^{I_0} \dots \partial_{\sigma_q}^{I_q} A_{i_1, \dots, i_q}$$

then the preceding equation (7.30) can be compactly written as follows:

$$B_0 A_{\sigma_0, \dots, \sigma_q} = 0.$$

In fact, due to symmetry properties of the type (3.18), namely:

$$A_{\sigma_{P(1)}, \dots, \sigma_{P(s)}, i_{Q(1)}, \dots, i_{Q(q)}}^{I_{P(1)}, \dots, I_{P(s)}} = (-1)^{|Q|} A_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{I_1, \dots, I_s} \quad (7.31)$$

we have

$$B_\alpha A_{\sigma_0, \dots, \sigma_q} = 0 \quad (\alpha = 0, \dots, q). \quad (7.32)$$

As a consequence we obtain:

$$B_0 \dots B_q A_{\sigma_0, \dots, \sigma_q} = 0$$

and lemma 3.6 can be applied; it follows that A_{\dots} has the generic form

$$A_{\dots} = \sum_{\alpha=0}^q B_\alpha A_{\dots}^\alpha$$

for some tensors $A_{\dots}^\alpha \in \mathcal{H}_{q, r, \dots, r, r-1, r, \dots, r}$ where the entry $r-1$ is on the position α .

One can plug this relation into the initial relation (7.32) and a similar relation is obtained for the tensors A_{\dots}^α . By recurrence, one gets:

$$A_{\dots} = \sum_{\alpha_0, \dots, \alpha_s=0}^q B_{\alpha_0} \dots B_{\alpha_s} A_{\dots}^{\alpha_0, \dots, \alpha_s} \quad (s = 0, \dots, q)$$

with $A_{\dots}^{\alpha_0, \dots, \alpha_s}$ some tensors in $\mathcal{H}_{q-s-1, r_0, \dots, r_q}$. In particular, for $s = q$ we obtain that in fact: $A_{\dots} = 0$ i.e.

$$\partial_{\sigma_0}^{I_0} \dots \partial_{\sigma_q}^{I_q} A_{i_1, \dots, i_q} = 0. \quad (7.33)$$

In other words, the functions A_{i_1, \dots, i_q} are polynomials in y_I^σ ($|I| = r$) of maximal degree q . So, the generic form of these functions is:

$$A_{i_1, \dots, i_q} = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s|=r} C_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{I_1, \dots, I_s} y_{I_1}^{\sigma_1} \cdots y_{I_s}^{\sigma_s} \quad (7.34)$$

with $C_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{I_1, \dots, I_s}$ some smooth functions on V^r having symmetry properties of the type (7.31). Remark the fact that these functions do not depend on the variables y_I^σ ($|I| = r$) and this justifies the fact that they live on the chart V^{r-1} .

Now we insert this generic expression into the projectability condition (7.29) and we get, in the same way as before

$$\mathcal{S}_{i_0, \dots, i_q}^- \mathcal{S}_{I_1 p}^+ \delta_{i_0}^p C_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{I_1, \dots, I_s} = 0 \quad (s = 1, \dots, q-1) \quad (7.35)$$

or, in tensor notations:

$$B_1 C_{\sigma_1, \dots, \sigma_q} = 0 \quad (s = 1, \dots, q-1).$$

In fact, due to the symmetry properties one has from here:

$$B_\alpha C_{\sigma_1, \dots, \sigma_q} = 0 \quad (\alpha = 1, \dots, s; \quad s = 1, \dots, q-1). \quad (7.36)$$

Using an argument familiar by now we get from here:

$$C_{\sigma_1, \dots, \sigma_q} = B_1 \cdots B_s \tilde{C}_{\sigma_1, \dots, \sigma_q} \quad (s = 1, \dots, q-1).$$

In full index notations this means that we have the following generic expression:

$$C_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{J_1 p_1, \dots, J_s p_s} = \mathcal{S}_{J_1 p_1}^+ \cdots \mathcal{S}_{J_s p_s}^+ \mathcal{S}_{i_1, \dots, i_q}^- \delta_{i_1}^{p_1} \cdots \delta_{i_s}^{p_s} A_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{J_1, \dots, J_s} \quad (s = 1, \dots, q) \quad (7.37)$$

where $|J_1| = \cdots = |J_s| = r-1$ and $A_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{J_1, \dots, J_s}$ are some smooth function on V^{r-1} completely antisymmetric in the indices i_1, \dots, i_q .

Inserting this expression in (7.34) one immediately obtains

$$A_{i_1, \dots, i_q} = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|J_1|, \dots, |J_s|=r-1} \mathcal{S}_{i_1, \dots, i_q}^- A_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{J_1, \dots, J_s} y_{J_1}^{\sigma_1} \cdots y_{J_s}^{\sigma_s}. \quad (7.38)$$

Let us remark that the expression $\mathcal{S}_{i_1, \dots, i_q}^- y_{J_1}^{\sigma_1} \cdots y_{J_s}^{\sigma_s}$ is completely antisymmetric in the couples $(I_1, \sigma_1), \dots, (I_s, \sigma_s)$ so one can consider that the tensors $A_{\sigma_1, \dots, \sigma_s, i_1, \dots, i_q}^{J_1, \dots, J_s}$ have the same property. In the end, it follows that they have the symmetry property (3.18).

The expression for the form ν becomes

$$\nu = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|J_1|, \dots, |J_s|=r-1} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{J_1, \dots, J_s} y_{J_1}^{\sigma_1} \cdots y_{J_s}^{\sigma_s} dx^{i_1} \wedge \cdots \wedge dx^{i_q} = h \tilde{\nu}_V \quad (7.39)$$

where

$$\tilde{\nu}_V \equiv \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|J_1|, \dots, |J_s|=r-1} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{J_1, \dots, J_s} dy_{J_1}^{\sigma_1} \wedge \cdots \wedge dy_{J_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \cdots \wedge dx^{i_q}. \quad (7.40)$$

This finishes the proof if we take into account that the relation (7.29) is equivalent to the projectability condition. ■

Now we are ready to obtain the most general form of a variationally trivial Lagrangian. First we note that we have

Proposition 7.3 *The Lagrange form $\lambda \in \Omega_{n,X}^r$ is variationally trivial iff on has:*

$$E_n([\lambda]) = 0. \quad (7.41)$$

Proof: It is an immediate consequence of the definition of a variationally trivial Lagrange form and of theorem 6.4. ■

The central result now follows:

Theorem 7.4 *Let $\lambda \in \Omega_{n,X}^r$ be a variationally trivial Lagrange form. Then for every point $j_x^r \gamma \in J^r Y$ there exists a neighbourhood V of $\gamma(x) \in Y$ and a n -form ρ_V defined in the chart (V^{r-1}, ψ^{r-1}) which is basic and we also have: (1) $\lambda = h\rho_V$ in the chart (V^r, ψ^r) ; (2) $d\rho_V = 0$. Conversely, if such a local form ρ_V exists, then the form λ is variationally trivial.*

Proof: (i) According to the previous proposition and applying the exactness of the variational sequence (theorem 6.3), there exists $\eta \in \Omega_{n-1}^r$ such that

$$[\lambda] = [d\eta],$$

or, equivalently

$$\lambda - d\eta \in \theta_n^r = \Omega_{n(c)}^r. \quad (7.42)$$

As a consequence we have

$$(\pi^{r+1,r})^* \lambda = h\lambda = hd\eta.$$

In particular, it follows that the n -form $hd\eta$ must be $\pi^{r+1,r}$ -projectable. We can apply proposition 7.1 and (7.42) rewrites as follows:

$$\lambda - d\nu \in \Omega_{n(c)}^r. \quad (7.43)$$

for some basic form ν . From here we get

$$h\lambda = hd\nu \quad (7.44)$$

or using (3.1)

$$(\pi^{r+1,r})^* \lambda = hd\nu. \quad (7.45)$$

Let us note that that this relation is completely equivalent to the initial condition of variationally triviality.

(ii) Next, one sees that from (7.44) it follows in particular that the n -form $d\nu$ is $\pi^{r+1,r}$ -projectable. We can apply proposition 7.2 and obtain that $\nu = h\tilde{\nu}_V$ for some basic form in the chart (V^{r-1}, ψ^{r-1}) . If we define

$$\rho_V \equiv d\tilde{\nu}_V \quad (7.46)$$

then we obtain after some computation that

$$(\pi^{r+1,r})^* (\lambda - h\rho_V) = 0$$

and (1) from the statement follows. The definition (7.46) guarantees that we also have (2). ■

Remark 7.4.1 A statement of the type appearing in this theorem is, in fact, valid for every $\lambda \in \Omega_{q,X}^r$ ($q \leq n$) such that $E_q([\lambda]) = 0$.

The theorem we just have proved has the following consequence (see also [3] thm. 4.3)

Corollary 7.5 Any variationally trivial Lagrange form $\lambda \in \Omega_{n,X}^r$ can be locally written as a total exterior derivative of a local form $\omega_V \in \mathcal{J}_{n-1}^r$:

$$\lambda = D\omega_V. \quad (7.47)$$

Proof: In the proof of the preceding theorem we restrict, eventually, the chart V^{r-1} and we have $\rho_V = d\eta_V$ for some $(n-1)$ -form on V^{r-1} . Then, according to the definition (3.66) of the total exterior derivative we have the formula from the statement with $\omega_V \equiv h\eta_V \in \mathcal{J}_{n-1}^r$. ■

One can now obtain the most general form of a variationally trivial local Lagrangian.

Theorem 7.6 Any variationally trivial local Lagrangian of order r has the following form in the chart (V^r, ψ^r) :

$$L = \sum_{s=0}^n \frac{1}{s!(n-s)!} \sum_{|I_1|, \dots, |I_s|=r-1} \mathcal{L}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} \mathcal{J}_{I_1, \dots, I_s}^{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}. \quad (7.48)$$

Here we have defined

$$\mathcal{J}_{I_1, \dots, I_s}^{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n} \equiv \varepsilon^{i_1, \dots, i_n} \prod_{l=1}^s y_{I_l i_l}^{\sigma_l} \quad (s = 0, \dots, n) \quad (7.49)$$

and the function \mathcal{L}_{\dots} are given by

$$\mathcal{L}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} \equiv \sum_{k=1}^s (-1)^{k-1} \partial_{\sigma_k}^{I_k} A_{\sigma_1, \dots, \sigma_k, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, \hat{I}_k, \dots, I_s} + \sum_{k=s+1}^q (-1)^{k-1} d_{i_k} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, \hat{i}_k, \dots, i_n}^{I_1, \dots, I_s}; \quad (7.50)$$

the expressions $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{n-1}}^{I_1, \dots, I_s}$ are smooth functions on V^{r-1} verifying the symmetry property (3.18) and $d_j = d_j^{r-1}$ is the corresponding formal derivative on V^{r-1} .

Proof: It is convenient to introduce the following forms:

$$\chi \equiv \sum_{s=0}^{n-1} \frac{1}{s!(n-1-s)!} \sum_{|I_1|, \dots, |I_s|=r-1} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{n-1}}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_{n-1}}$$

and

$$\theta \equiv \sum_{s=0}^n \frac{1}{s!(n-s)!} \sum_{|I_1|, \dots, |I_s|=r-1} \mathcal{L}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_n}.$$

Then one finds out by direct computation that

$$\theta = d\chi + \text{contact terms.}$$

Next, one takes in theorem 7.4 $\tilde{\nu}_V = \chi$ and it follows that $\rho_V = \theta + \text{contact terms}$. Finally, by direct computation one discovers that $\lambda = h\rho_V$ has the expression (7.48). ■

Remark 7.6.1 *Let us note that the expressions (7.50) are of the same type as those given by (7.5).*

The expressions (7.49) defined above are called *hyper-Jacobians* [7] [15] (see these references for similar results). It is immediate that they have antisymmetry properties of the type (3.18).

Now we give another argument for the converse statement from the preceding theorem is true. First we have:

Corollary 7.7 *The local expression of a variationally trivial Lagrangian (7.48) can be rewritten as follows:*

$$L = d_j V^j \quad (7.51)$$

where V^j are some smooth functions on V^r .

Proof: Using the notations introduced in the proof above let us define the local expressions on V^r :

$$V^j \equiv \varepsilon^{j,i_1,\dots,i_{n-1}} \sum_{s=0}^n \frac{1}{s!(n-1-s)!} \sum_{|I_1|,\dots,|I_s|=r-1} A_{\sigma_1,\dots,\sigma_s,i_{s+1},\dots,i_{n-1}}^{I_1,\dots,I_s} y_{I_1 i_1}^{\sigma_1} \cdots y_{I_s i_s}^{\sigma_s}.$$

One checks now that the formula from the statement is true. ■

Now we indeed have:

Theorem 7.8 *The expression (7.48) is variationally trivial.*

Proof: We have according to the preceding corollary $E_\sigma(L) = E_\sigma(d_j V^j) = 0$ because we can apply (5.5). ■

Remark 7.8.1 *Some globalisation of the results above can be found in [22] (see theorem 5 and corollary 1 from this reference). In particular, for $r = 1$ one obtains known results [25], [8], [16], [14].*

We close this Section with the analysis of the following problem. We have discovered that a variationally trivial Lagrangian depends on the highest order derivatives through some very particular polynomial expressions. The problem is to obtain a system of partial differential equations which is compatible only with this structure. More precisely, we have the following result.

Theorem 7.9 *Let us suppose that the local Lagrangian L on V^r verifies the system of partial differential equations:*

$$\mathcal{S}_{p_1,\dots,p_r,j_r}^+ \partial_\rho^{p_1,\dots,p_r} \partial_\sigma^{j_1,\dots,j_r} L = 0. \quad (7.52)$$

Then L is a polynomial in y_I^σ , $|I| = r$ of the following form:

$$L = \sum_{s=0}^n \frac{1}{s!(n-s)!} \sum_{|I_1|,\dots,|I_s|=r-1} \mathcal{L}_{\sigma_1,\dots,\sigma_s,i_{s+1},\dots,i_n}^{I_1,\dots,I_s} \mathcal{J}_{I_1,\dots,I_s}^{\sigma_1,\dots,\sigma_s,i_{s+1},\dots,i_n} \quad (7.53)$$

where $\mathcal{L}_{\sigma_1,\dots,\sigma_s,i_{s+1},\dots,i_n}^{I_1,\dots,I_s}$ are some smooth functions on V^{r-1} verifying symmetry properties of the type (3.18). Conversely, if L is of the form above, then the system (7.52) is identically fulfilled.

Proof:

(i) We will first prove that if the local Lagrangian L verifies the identities:

$$\mathcal{S}_{p_1, \dots, p_{r-l+1}, j_{k-l+1}, \dots, j_r}^+ \partial_{\rho}^{p_1, \dots, p_{r-l+1}} \partial_{\sigma}^{j_1, \dots, j_k} L = 0 \quad (1 \leq l \leq k \leq r) \quad (7.54)$$

and

$$\sum_{k=0}^r (-1)^k d_{j_1}^r \cdots d_{j_k}^r \partial_{\sigma}^{j_1, \dots, j_k} L = 0 \quad (7.55)$$

then we have $E_{\sigma}(L) = 0$ in V^s ($s > 2r$).

Indeed one starts directly from the definition

$$E_{\sigma}(L) = \sum_{k=0}^r (-1)^k d_{j_k}^s \cdots d_{j_1}^s \partial_{\sigma}^{j_1, \dots, j_k} L = \sum_{k=0}^r (-1)^k (d_{j_k}^r + \sum_{l=r}^{2r-1} y_{p_1, \dots, p_l, k_k}^{\nu} \partial_{\nu}^{p_1, \dots, p_l}) d_{j_{k-1}}^s \cdots d_{j_1}^s \partial_{\sigma}^{j_1, \dots, j_k} L \quad (7.56)$$

and commutes $\partial_{\nu}^{p_1, \dots, p_l}$ over $d_{j_{k-1}}^s \cdots d_{j_1}^s$.

We must use the hypothesis (7.54) and a generalization of the formula (2.23), namely

$$[\partial_{\sigma}^{p_1, \dots, p_k}, d_{j_1} \cdots d_{j_l}] = \mathcal{S}_{p_1, \dots, p_k}^+ \mathcal{S}_{j_1, \dots, j_l}^+ \sum_{t=1}^l c_{k, l, t} \delta_{j_1}^{p_1} \cdots \delta_{j_t}^{p_t} d_{j_{t+1}} \cdots d_{j_l} \partial_{\sigma}^{p_{t+1}, \dots, p_k} \quad (k \geq l) \quad (7.57)$$

where $c_{k, l, t} \in \mathbb{R}_+$; this formula can be proved by induction on l .

Now we easily obtain that in fact the terms corresponding to the sum over l in (7.56) give a null contribution, so we are left with:

$$E_{\sigma}(L) = \sum_{k=0}^r (-1)^k d_{j_k}^r d_{j_{k-1}}^s \cdots d_{j_1}^s \partial_{\sigma}^{j_1, \dots, j_k} L.$$

We continue in the same way by recurrence and finally get

$$E_{\sigma}(L) = \sum_{k=0}^r (-1)^k d_{j_1}^r \cdots d_{j_k}^r \partial_{\sigma}^{j_1, \dots, j_k} L$$

which is zero, according to (7.55).

It appears that (7.54) and (7.55) imply that the Lagrangian L is variationally trivial. According to theorem 7.4, L has the form (7.53) from the statement, but with some restrictions on the functions \mathcal{L}_{\dots} . On the other hand, it is clear that the dependence on the highest order derivatives must follow only from the equations containing only the partial derivatives of order r i.e. (7.54) for $k = r$ and $l = 1$ i.e. the system (7.52) from the statement.

(ii) The converse statement follows rather easily. One shows by direct computations that if L is given by the formula (7.53), then:

$$\partial_{\sigma_1}^{I_1} L = \sum_{s=1}^n (-1)^{s-1} \frac{1}{(s-1)!(n-s)!} \sum_{|I_2|, \dots, |I_s|=r-1} (B_1 \mathcal{L})_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} \mathcal{J}_{I_1, \dots, I_s}^{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}.$$

Iterating the derivation procedure we have

$$\partial_{\sigma_1}^{I_1} \partial_{\sigma_2}^{I_2} L = - \sum_{s=2}^n \frac{1}{(s-2)!(n-s)!} \sum_{|I_3|, \dots, |I_s|=r-1} (B_1 B_2 \mathcal{L})_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} \mathcal{J}_{I_1, \dots, I_s}^{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}$$

and one proves that (7.52) is fulfilled by direct computation. ■

8 Locally Variational Differential Equations

The definitions for a general differential equation and for a locally variational differential equation have been given previously (see the formulæ (3.47) and resp. (5.22).) We want to analyse the general structure of a locally variational differential equation along the same lines of argument as in the previous Section. We will not be able to obtain the most general expression for such an object (as we have been able to obtain in the case of variationally trivial Lagrangians) but we will produce a generic expression of the same type as (7.53). We mention again that this result has been already obtained in [2] with a completely different method.

Our starting point is a analogue of the proposition 7.1 for the case $q = n$; it is natural to also make the replacement $h \rightarrow p_1$.

Proposition 8.1 *Let $\eta \in \Omega_n^r$ such that $p_1 d\eta$ is a $\pi^{r+1,r}$ -projectable $(n+1)$ -form. Then one can write η as follows:*

$$\eta = \nu + \sum_{|J| \leq r-1} \Phi_{\sigma, i_1, \dots, i_{n-1}}^J \omega_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_{n-1}} + d\phi + \psi \quad (8.1)$$

where $\nu \in \Omega_{n,X}^r$ is a basic n -form, $\psi \in \Omega_{n+1}^r$ is a 2-contact form and $\Phi_{\sigma, i_1, \dots, i_{n-1}}^J$ are smooth functions on V^r completely antisymmetric in the indices i_1, \dots, i_{n-1} .

Proof:

We proceed in analogy with proposition 7.1. Let (V, ψ) be a chart on Y . Then in the associated chart (V^r, ψ^r) one can write η in the standard form

$$\eta = \eta_0 + \eta_1 \quad (8.2)$$

where in η_0 we collect all terms containing at least one factor ω_J^σ ($|J| \leq r-1$) and η_1 is a polynomial only in dx^i and dy_J^σ ($|J| = r+1$):

$$\eta_1 = \sum_{s=0}^n \sum_{|I_1|, \dots, |I_s|=r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_n}, \quad (8.3)$$

where the coefficients $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s}$ have antisymmetry properties of type (3.18).

The generic form of η_0 is

$$\eta_0 = \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J \quad (8.4)$$

with Φ_σ^J some $(n-1)$ -forms. It follows then that

$$d\eta = d\eta_1 + \sum_{|J| \leq r-1} \left[(d\omega_J^\sigma) \wedge \Phi_\sigma^J - \omega_J^\sigma \wedge (d\Phi_\sigma^J) \right]. \quad (8.5)$$

One applies the operator p_1 to this equality and uses (4.1); a new relation is obtained, namely:

$$p_1 d\eta = p_1 d\eta_1 + \sum_{|J| \leq r-1} \left[(d\omega_J^\sigma) \wedge h\Phi_\sigma^J - \omega_J^\sigma \wedge (hd\Phi_\sigma^J) \right]. \quad (8.6)$$

The expression of $d\eta_1$ has been computed quite generally before and is given by (7.5). Applying the operator p_1 and using (4.4) one obtains:

$$p_1 d\eta_1 = \sum_{s=0}^n \sum_{|I_1|, \dots, |I_s|=r} \sum_{|J| \leq r-1} \left(\partial_\nu^J A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} \right) y_{I_1}^{\sigma_1} \cdots y_{I_s}^{\sigma_s} \omega_J^\nu \wedge dx^{i_{s+1}} \wedge \cdots \wedge dx^{i_n} \\ + \sum_{|I_1|=r} \tilde{B}_{\sigma_1, i_2, \dots, i_{n+1}}^{I_1} \omega_{I_1}^{\sigma_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_{n+1}},$$

where

$$\tilde{B}_{\sigma_1, i_2, \dots, i_{n+1}}^{I_1} = \mathcal{A}_{i_2, \dots, i_{n+1}} \sum_{s=1}^{n+1} s \sum_{|I_1|, \dots, |I_s|=r} \tilde{A}_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{n+1}}^{I_1, \dots, I_s} y_{I_2}^{\sigma_2} \cdots y_{I_s}^{\sigma_s}. \quad (8.7)$$

The expression (8.6) becomes

$$p_1 d\eta = - \sum_{|J|=r-1} \omega_{Ji}^\sigma \wedge dx^i \wedge h\Phi_\sigma^J + \sum_{|I_1|=r} \tilde{B}_{\sigma_1, i_2, \dots, i_{n+1}}^{I_1} \omega_{I_1}^{\sigma_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_{n+1}} + \cdots, \quad (8.8)$$

where by \cdots we mean contributions which do not contain the differentials ω_I^σ , $|I| = r$.

The first term has the generic form

$$\sum_{|I_1|=r} C_{\sigma_1, i_2, \dots, i_{n+1}}^{I_1} \omega_{I_1}^{\sigma_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_{n+1}}$$

where $C_{\sigma_1, i_2, \dots, i_{n+1}}^{I_1}$ are some polynomials in y_I^σ , $|I| = r+1$ of maximal degree $(n-1)$; because of the presence of the combination $\omega_{Ji}^\sigma \wedge dx^i$ these polynomials are of delta-type i.e. they are obtained by applying B_1 on some other tensors; it follows that we have in compact tensor notations:

$$B_1 C_\sigma = 0. \quad (8.9)$$

The expression (8.8) does not depend on y_I^σ , $|I| = r+1$ by hypothesis. This is equivalent with the independence of $\tilde{B}_\sigma - C_\sigma$ on y_I^σ , $|I| = r+1$. In particular, the same thing must be true for $B_1(\tilde{B}_\sigma - C_\sigma) = B_1 \tilde{B}_\sigma$ where use of (8.9) has been made.

If the expression (8.7) is used one obtains

$$\mathcal{S}_{i_2, \dots, i_{n+1}}^-(B_1 \tilde{A})_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{n+1}}^{I_1, \dots, I_s} y_{I_2}^{\sigma_2} \cdots y_{I_s}^{\sigma_s} = 0, \quad (s = 2, \dots, n+1)$$

or, in compact tensor notations:

$$B_2 \cdots B_s B_1 \tilde{A}_{\sigma_1, \dots, \sigma_s} = 0, \quad (s = 2, \dots, n+1).$$

One can apply lemma 3.6 and obtains that $\tilde{A}_{\sigma_1, \dots, \sigma_s}$ has the following form

$$\tilde{A}_{\sigma_1, \dots, \sigma_s} = \sum_{\alpha=1}^s B_\alpha A_{\sigma_1, \dots, \sigma_s}^\alpha, \quad (s = 2, \dots, n+1).$$

This expression must be substituted into the formula for $d\eta_1$ and, after some elementary prelucrations, the following generic form is produced:

$$d\eta_1 = \nu_1 + d\phi + \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J. \quad (8.10)$$

Here $\phi \in \Omega_{n(c)}^r$ is a contact form, ν_1 has the form

$$\nu_1 = \sum_{|I_1|=r} A_{\sigma_1, i_2, \dots, i_{n+1}}^{I_1} dy_{I_1}^{\sigma_1} \wedge dx^{i_2} \wedge \dots \wedge dx^{i_{n+1}} \quad (8.11)$$

and Φ_σ^J is a polynomial of degree n in dx^i and dy_I^σ , $|I| = r$. Explicitly:

$$\Phi_\sigma^J = \sum_{s=0}^n \sum_{|I_1|, \dots, |I_s|=r} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{J, I_1, \dots, I_s} dy_{I_1}^{\nu_1} \wedge \dots \wedge dy_{I_s}^{\nu_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_n}; \quad (8.12)$$

here we can assume a (partial) symmetry property of the type (7.12) and we make the convention

$$\Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{J, I_1, \dots, I_s} = 0, \quad \forall J \text{ s.t. } |J| \geq r. \quad (8.13)$$

(ii) From the expression (8.10) one obtains after exterior differentiation and application of the operator p_2 the following condition:

$$p_2 d\nu_1 - \sum_{|J| \leq r-1} (\omega_{Ji}^\sigma \wedge dx^i \wedge p_1 \Phi_\sigma^J - \omega_J^\sigma \wedge p_1 d\Phi_\sigma^J) = 0. \quad (8.14)$$

As in the proof of the proposition 7.1, one finds out from this equation the following consequences:

$$\mathcal{S}_{i_2, \dots, i_{n+1}}^- \mathcal{S}_{I_1 p_1}^+ \dots \mathcal{S}_{I_s p_s}^+ \left[\tilde{\Phi}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_{n+1}}^{\{j_1, \dots, j_k\}, I_1, \dots, I_s} + (-1)^s \mathcal{S}_{j_1, \dots, j_k} \delta_{i_{s+1}}^{j_1} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+2}, \dots, i_{n+1}}^{\{j_2, \dots, j_k\}, I_1, \dots, I_s} \right] \times \\ \delta_{i_2}^{p_2} \dots \delta_{i_s}^{p_s} = 0, \quad (s = 2, \dots, n+1, k = 0, \dots, r-1) \quad (8.15)$$

and

$$\mathcal{S}_{i_2, \dots, i_{n+1}}^- \mathcal{S}_{I_1 p_1}^+ \dots \mathcal{S}_{I_s p_s}^+ \mathcal{S}_{j_1, \dots, j_k}^+ \delta_{i_{s+1}}^{j_1} \left(\Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+2}, \dots, i_{n+1}}^{\{j_2, \dots, j_r\}, I_1, \dots, I_s} + \Phi_{\nu_1, \sigma, \nu_2, \dots, \nu_s, i_{s+2}, \dots, i_{n+1}}^{I_1, \{j_2, \dots, j_k\}, I_2, \dots, I_s} \right) \times \\ \delta_{i_2}^{p_2} \dots \delta_{i_s}^{p_s} = 0, \quad (s = 2, \dots, n+1). \quad (8.16)$$

(Here use have been made of the definition (7.15).)

In compact notations, these two relations can be written in a similar way and with similar notations (using, in particular, the convenient operator (7.16)):

$$B_2 \dots B_s \left((\Delta' \Phi)_s^k + B_0 \Phi_s^{k-1} \right) = 0, \quad (s = 2, \dots, n+1, k = 0, \dots, r-1) \quad (8.17)$$

and

$$B_2 \dots B_s \left(B_0 \Phi_s^{r-1} + B_1 \Psi_s^{r-1} \right) = 0, \quad (s = 2, \dots, n+1); \quad (8.18)$$

here we have defined:

$$\Psi_{\nu_0, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_{n+1}}^{I_0, I_1, \dots, I_s} \equiv \Phi_{\nu_1, \nu_0, \dots, \nu_s, i_{s+1}, \dots, i_{n+1}}^{I_1, I_0, \dots, I_s}. \quad (8.19)$$

It is clear that we must apply again the descent procedure from the proof of the proposition 7.1 in a case which is a little more complicated. We first deal with the equation (8.18) by applying the operator B_1 ; then one gets:

$$B_0 \dots B_s \Phi_s^{r-1} = 0, \quad (s = 2, \dots, n+1)$$

from where one obtains, with lemma 3.6

$$\Phi_s^{r-1} = \sum_{\alpha=0}^s B_\alpha \Phi_s^{r-1, \alpha}. \quad (s = 2, \dots, n+1).$$

As before, one can redefine Φ_s^{r-1} such that, instead of the previous formula we have a generic expression of the type:

$$\Phi_s^{r-1} = B_0 C_s^{r-1}, \quad (s = 2, \dots, n+1) \quad (8.20)$$

and, instead of (8.10) we have

$$d\eta_1 = \nu_1 + d\phi + \sum_{|J| \leq r-1} \omega_J^\sigma \wedge \Phi_\sigma^J + \psi \quad (8.21)$$

where ψ is a 2-contact form.

One can take from the beginning the previous argument, based on acting on this relation with the operator $p_2 d$, and obtains this time only the relation (8.17) together with (8.20); the tensors C_s^{r-1} stay arbitrary. Now we are back in the same case we have previously analysed in proposition 7.1. The descent procedure can be applied and instead of (8.21) one gets the following expression:

$$\begin{aligned} d\eta_1 = \nu_1 + d\phi + \psi + \sum_{|J| \leq r-1} \Phi_{\sigma, i_1, \dots, i_n}^J \omega_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_n} + \\ \sum_{|J| \leq r-1} \sum_{|I_1|=r} \Phi_{\sigma, \nu_1, i_2, \dots, i_n}^{J I_1} \omega_J^\sigma \wedge dy_{I_1}^{\nu_1} \wedge dx^{i_2} \wedge \dots \wedge dx^{i_n} \end{aligned} \quad (8.22)$$

where the notations have the same meaning as before.

Now we apply for the third time the argument based on acting on this relation with the operator $p_2 d$, and obtain instead of (8.17) and (8.18) only the equation

$$B_2(\Delta' \Phi^k)_2 = 0, \quad (k = 0, \dots, r-1).$$

From symmetry considerations we also have the same relation with $B_2 \rightarrow B_1$ so, we can obtain as in the proof of proposition 7.2, that the following equality stays true:

$$\Delta' \Phi_1^k = B_1 B_2 C^k, \quad (k = 0, \dots, r-1) \quad (8.23)$$

where $C^k \in \mathcal{H}_{n-1, k-2, r-1, r-1}$.

(iii) We concentrate on the analysis of the previous equation (8.23). The key observation is that there exists a homotopy operator for Δ' . Indeed, we have

Lemma 8.2 *Suppose that the tensor $A = \oplus_{s=1}^q A_s$ verifies the equation:*

$$\Delta' A = 0. \quad (8.24)$$

Then there exists a tensor $C = \oplus_{s=0}^{q-1} C_s$ such that

$$A = \Delta' C. \quad (8.25)$$

Proof:

An explicit formula for the tensor C (suggested by [22]) is the following one. One defines the map $\chi : \mathbb{R} \times V^r \rightarrow V^r$ by

$$\chi(u, (x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_r}^\sigma)) = (x^i, y^\sigma, y_j^\sigma, \dots, u y_{j_1, \dots, j_r}^\sigma)$$

and afterwards:

$$C_{\nu, \sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q-1}}^{J, I_1, \dots, I_s} \equiv (s+1) \sum_{|I_0|=r} y_{I_0}^{\sigma_0} \int_0^1 u^s A_{\nu, \sigma_0, \dots, \sigma_s, i_{s+1}, \dots, i_{q-1}}^{J, I_0, \dots, I_s} \circ \chi \quad du, \quad (s = 1, \dots, q-1).$$

By elementary computations one finds out that:

$$(\Delta' C)_{\nu, \sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{J, I_1, \dots, I_s} = A_{\nu, \sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{J, I_1, \dots, I_s}, \quad (s = 1, \dots, q)$$

and this gives us the desired homotopy formula. ∇

The previous lemma has a generalization.

Lemma 8.3 *Let us suppose that we have the equation:*

$$\Delta' A = D. \quad (8.26)$$

Then tensor D verifies the consistency relation:

$$\Delta' D = 0 \quad (8.27)$$

and a particular solution of the equation above is of the form:

$$A_{\nu, \sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_{q-1}}^{J, I_1, \dots, I_s} = (s+1) \sum_{|J|=r} y_{I_0}^{\sigma_0} \int_0^1 u^s D_{\nu, \sigma_0, \dots, \sigma_s, i_{s+1}, \dots, i_{q-1}}^{J, I_0, \dots, I_s} \circ \chi \quad du, \quad (s = 1, \dots, q-1). \quad (8.28)$$

Proof: The consistency condition follows from (7.20) and the last equality by direct computations. ∇

(iv) We need the preceding two lemmas only in the case $q = n$. One finds out that the generic solution of the equation (8.23) is

$$\Phi_1^k = \Delta' \Psi^k + B_1 A^k, \quad (k = 0, \dots, r-1) \quad (8.29)$$

where the first term is an arbitrary solution of the homogeneous equation (obtained with the first lemma) and the second term is a particular solution of the non-homogeneous equation (obtained with the second lemma for $D = B_1 B_2 C^k$.)

Now we must substitute this result in the last sum from the formula (8.22); it is not very hard to regroup the result with the first sum such that one obtains a more simple expression:

$$d\eta_1 = \nu_1 + d\phi + \psi + \sum_{|J| \leq r} A_{\sigma, i_1, \dots, i_n}^J dy_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_n}. \quad (8.30)$$

(v) The formula above can be prelucrated, as in proposition 7.1, using the contact homotopy operator A . Indeed, if we apply to the form $\eta_1 - \phi_1$ the formula (6.7) we obtain:

$$\eta_1 - \phi_1 = A \left(\nu_1 + \sum_{|J| \leq r} A_{\sigma, i_1, \dots, i_n}^J dy_J^\sigma \wedge dx^{i_1} \wedge \dots \wedge dx^{i_n} + \psi \right) + dA(\eta_1 - \phi_1) + \eta_1(0) - \phi_1(0). \quad (8.31)$$

As before we easily establish the following facts:

- $A\nu_1 = 0$;
- $A(\sum \dots)$ and $\eta_1(0)$ are basic n -forms;
- if ϕ is contact, then $\phi_1(0)$ is contact;
- if ψ is a 2-contact form, then $A\psi$ is a contact form.

These results, gives us the following generic expression for η_1 :

$$\eta_1 = \nu + \eta_0 + d\phi \quad (8.32)$$

where ν is a basic n -form, ϕ is a contact $(n-1)$ -form and η_0 is a n -contact form of the type (8.4). It is obvious that the same type of expression stays true for η also.

(vi) It is useful to notice that the forms Φ_σ^J from (8.4) can also be decomposed into a contribution having at least a factor ω_J^σ and a polynomial of degree $(n-1)$ in the differentials dx^i and dy_I^σ , $|I| = r$. In this way, the formula for η takes a more convenient form, namely:

$$\eta = \nu + \eta_0 + d\phi + \psi \quad (8.33)$$

where ν and ϕ have the same properties as above, ψ is a 2-contact form and η_0 has the expression of the type (8.4) but the forms Φ_σ^J are polynomials (of degree $n-1$) only in the differentials dx^i and dy_I^σ , $|I| = r$. Explicitly:

$$\Phi_\sigma^J = \sum_{s=0}^{n-1} \sum_{|I_1|, \dots, |I_s|=r} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_{n-1}}^{J, I_1, \dots, I_s} dy_{I_1}^{\nu_1} \wedge \dots \wedge dy_{I_s}^{\nu_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_{n-1}} \quad (8.34)$$

where symmetry properties of the type (7.12) can be imposed and we admit that:

$$\Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{J, I_1, \dots, I_s} = 0, \quad \forall J \quad s.t. \quad |J| \geq r. \quad (8.35)$$

We must impose on this expression for η the condition from the statement of the proposition. The local expression for ν is given by (2.32). We obtain instead of (8.6):

$$p_1 d\eta = \sum_{|J| \leq r} (\partial_\sigma^J L) \omega_J^\sigma \wedge \theta_0 + \sum_{|J| \leq r-1} \left[(d\omega_J^\sigma) \wedge h\Phi_\sigma^J - \omega_J^\sigma \wedge (hd\Phi_\sigma^J) \right]. \quad (8.36)$$

The condition of $\pi^{r+1, r}$ -projectability for this expression amounts to a set of relation similar to (8.15), namely:

$$\mathcal{S}_{i_1, \dots, i_n}^- \mathcal{S}_{I_1 p_1}^+ \dots \mathcal{S}_{I_s p_s}^+ \left[\tilde{\Phi}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{j_1, \dots, j_k, I_1, \dots, I_s} + (-1)^s \mathcal{S}_{j_1, \dots, j_k} \delta_{i_{s+1}}^{j_1} \Phi_{\sigma, \nu_1, \dots, \nu_s, i_{s+2}, \dots, i_n}^{j_2, \dots, j_k, I_1, \dots, I_s} \right] \times \\ \delta_{i_1}^{p_1} \dots \delta_{i_s}^{p_s} = 0, \quad (s = 1, \dots, n, k = 0, \dots, r) \quad (8.37)$$

In compact notations this amounts to

$$B_1 \cdots B_s \left((\Delta' \Phi)_s^k + B_0 \Phi_s^{k-1} \right) = 0, \quad (s = 1, \dots, n, k = 0, \dots, r) \quad (8.38)$$

which can be analysed with the, by now familiar, descent technique. Some terms can be grouped into the exterior differential of a contact form and in this way the formula from the statement emerges. ■

The converse of this proposition is not true. We proceed nevertheless to the study of locally variational differential equations by proving an analogue of proposition 7, namely:

Proposition 8.4 *Let T be a differential equation. Then T is locally variational iff*

$$E_{n+1}([T]) = 0. \quad (8.39)$$

Proof: It is an immediate consequence of the definition of a variationally trivial Lagrange form and of theorem 6.5. ■

Then we have the central result of this section:

Theorem 8.5 *Let T be a locally variational differential equation. Then T has the following local form in V^r given by (3.47) where:*

$$T_\sigma = \varepsilon^{i_1, \dots, i_n} \sum_{s=0}^n \sum_{|I_1|, \dots, |I_s|=r-1} \mathcal{T}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} \mathcal{J}_{I_1, \dots, I_s}^{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_n} \quad (8.40)$$

where $\mathcal{T}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s}$ are some smooth functions on V^{r-1} verifying symmetry properties of the type (3.18); this symmetry property leaves aside the index σ .

Proof: (i) According to the previous proposition and applying the exactness of the variational sequence (theorem 6.3), there exists $\lambda \in \Omega_n^r$ such that

$$[T] = [d\lambda],$$

or, equivalently

$$T - d\lambda \in \theta_{n+1}^r.$$

If we use (6.1) we obtain that

$$T - d\lambda = \alpha + d\beta$$

where $\alpha \in \Omega_{n+1(c)}^r$ and $\beta \in \Omega_{n(c)}^r$. We can redefine $\lambda \rightarrow \lambda - \beta$ and we obtain

$$T - d\lambda = \alpha \in \Omega_{n+1(c)}^r \quad (8.41)$$

or, equivalently:

$$p_1 T = p_1 d\lambda. \quad (8.42)$$

(ii) In particular, it follows that the $(n+1)$ -form $p_1 d\lambda$ must be $\pi^{r+1, r}$ -projectable. We can apply proposition 8.1 and obtain that λ has the following expression:

$$\lambda = \nu + \sum_{|J| \leq r-1} \Phi_{\sigma, i_1, \dots, i_{n-1}}^J \omega_J^\sigma \wedge dx^{i_1} \wedge \cdots \wedge dx^{i_{n-1}} + d\phi + \psi \quad (8.43)$$

where $\nu \in \Omega_{n,X}^r$ is a basic n -form, $\psi \in \Omega_{n+1}^r$ is a 2-contact form and $\Phi_{\sigma,i_1,\dots,i_{n-1}}^J$ $|J| = 1, \dots, r-1$ are smooth functions on V^r completely antisymmetric in the indices i_1, \dots, i_{n-1} ; for uniformity of notations we will extend the sum to $|J| = r$ with the convention:

$$\Phi_{\sigma,i_1,\dots,i_{n-1}}^J = 0, \quad |J| = r. \quad (8.44)$$

One computes the exterior differential of this form and imposes the condition of projectability. As in the proposition 7.2 one finds out that $p_1 d\lambda$ is $\pi^{r+1,r}$ -projectable *iff* the following equations are fulfilled:

$$\mathcal{S}_{i_1,\dots,i_n}^- \delta_{i_1}^p \tilde{\Phi}_{\sigma,\nu_1,i_2,\dots,i_n}^{J,I_1} = 0 \quad (8.45)$$

or :

$$\mathcal{S}_{i_1,\dots,i_n}^- \mathcal{S}_{Ip} \delta_{i_1}^p \left(\partial_\sigma^I \Phi_{\sigma,i_2,\dots,i_n}^J \right) = 0. \quad (8.46)$$

This equation is of the same type as (7.30) and the analysis performed there can be applied (one notices that the index q can take the value $n-1$). As a result, Φ_σ^J has a polynomial of the type (7.38)

$$\Phi_{\sigma,i_1,\dots,i_{n-1}}^J = \sum_{s=0}^{n-1} \sum_{|I_1|,\dots,|I_s|=r-1} \mathcal{S}_{i_1,\dots,i_{n-1}}^- C_{\sigma,\nu_1,\dots,\nu_s,i_1,\dots,i_{n-1}}^{J,I_1,\dots,I_s} y_{I_1 i_1}^{\nu_1} \cdots y_{I_s i_s}^{\nu_s} \quad (8.47)$$

where one can suppose that the tensors $C_{\sigma,\nu_1,\dots,\nu_s,i_1,\dots,i_{n-1}}^{J,I_1,\dots,I_s}$ have (partial) symmetry properties of the type (7.12).

(iii) Now one inserts the expressions (3.47) and (8.43) for T and respectively for λ into the projectability condition (8.42) and obtains after some algebra the following two relations:

$$T_\sigma = \partial_\sigma L + \varepsilon^{i_1,\dots,i_n} \tilde{\Phi}_{\sigma,i_1,\dots,i_n}^\emptyset \quad (8.48)$$

and

$$\partial_\sigma^J L = \varepsilon^{i_1,\dots,i_n} \left[\sum_{s=0}^{n-1} \sum_{|I_1|,\dots,|I_s|=r-1} (B_0 C)_{\sigma,\nu_1,\dots,\nu_s,i_{s+1},\dots,i_n}^{J,I_1,\dots,I_s} y_{I_1 i_1}^{\nu_1} \cdots y_{I_s i_s}^{\nu_s} + \tilde{\Phi}_{\sigma,i_1,\dots,i_n}^J \right], \quad (|J| = 1, \dots, r). \quad (8.49)$$

One must substitute into the last equation the expressions (8.47) for $\tilde{\Phi}_{\sigma,i_1,\dots,i_n}^J$ and after some computations one arrives at

$$\begin{aligned} \partial_\sigma^J L = F_\sigma^J \equiv \varepsilon^{i_1,\dots,i_n} \sum_{s=0}^{n-1} \sum_{|I_1|,\dots,|I_s|=r-1} \left[(B_0 C)_{\sigma,\nu_1,\dots,\nu_s,i_{s+1},\dots,i_n}^{J,I_1,\dots,I_s} + (\Delta' C)_{\sigma,\nu_1,\dots,\nu_s,i_{s+1},\dots,i_n}^{J,I_1,\dots,I_s} \right] \times \\ y_{I_1 i_1}^{\nu_1} \cdots y_{I_s i_s}^{\nu_s}, \quad (|J| = 1, \dots, r). \end{aligned} \quad (8.50)$$

It is clear that Frobenius conditions of integrability should be fulfilled:

$$\partial_\zeta^K F_\sigma^J = \partial_\sigma^J F_\zeta^K, \quad |J|, |K| = 1, \dots, r. \quad (8.51)$$

Instead of trying to solve directly these equations we proceed as follows. We admit that the integrability conditions (8.51) are fulfilled for $|J| = |K| = r$. Then one can obtain from (8.50) with a well-known homotopy formula:

$$L = L_0 + \int_0^1 \sum_{|J|=r} y_J^\sigma F_\sigma^J(x^i, y^\sigma, y_j^\sigma, \dots, y_{j_1, \dots, j_{r-1}}^\sigma, \dots, u y_{j_1, \dots, j_r}^\sigma) du \quad (8.52)$$

where L_0 does not depend on the highest order derivatives. Using the explicit expression for F_σ^J from (8.50) one arrives at:

$$L = L_0 + \varepsilon^{i_1, \dots, i_n} \sum_{s=1}^n \frac{1}{s} \sum_{|I_1|, \dots, |I_s|=r-1} A_{\nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} y_{I_1 i_1}^{\nu_1} \cdots y_{I_s i_s}^{\nu_s} \quad (8.53)$$

where

$$A_{\nu_0, \dots, \nu_s, i_{s+1}, \dots, i_{n-1}}^{I_0, \dots, I_s} \equiv \mathcal{S}_{(I_0, \nu_0), \dots, (I_s, \nu_s)}^- C_{\nu_0, \dots, \nu_s, i_{s+1}, \dots, i_{n-1}}^{I_0, \dots, I_s}, \quad (s = 0, \dots, n-1) \quad (8.54)$$

is constructed from C_{\dots} such that it verifies the complete symmetry property (3.18). Now one computes from (8.52) the partial derivatives of L and, considering the highest order ones, obtains by comparison with the relation (8.50)

$$B_0 \cdots B_s (C_s - A_s) = 0, \quad s = 0, \dots, n-1.$$

If we apply lemma 3.6 one obtains, as usual, that

$$C_{\nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} = A_{\nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s} + \delta - \text{terms}.$$

If we substitute this expression for C_{\dots} into the equation (8.47) for $|J| = r-1$ we obtain that the delta terms give a null contribution; as a consequence, it follows that the expressions $C_{\nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s}$, $|I_1|, \dots, |I_s| = r-1$ can be considered to have the symmetry property (3.18) without loosing the generality. Then the integrability condition (8.51) is fulfilled for $|J| = r$.

(iv) Finally, we develop the expression (8.48) for T_σ using the expression (8.47) for $\tilde{\Phi}_{\sigma, i_1, \dots, i_n}^\emptyset$ and the expression (8.52) for L . It is elementary to prove that both terms in (8.48) are polynomials in the hyper-Jacobians of the type appearing in the statement of the theorem. ■

Remark 8.5.1 The functions $\mathcal{T}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{I_1, \dots, I_s}$ appearing in the statement cannot be completely arbitrary because we did not use all the integrability conditions (8.51).

Remark 8.5.2 For the case $r = 2$ analysed in detail in [11] it is possible to use completely the Helmholtz-Sonin equations involving the highest order derivatives. One obtains that T_σ has a expression as in the statement of the theorem, but the coefficients $\mathcal{T}_{\sigma, \nu_1, \dots, \nu_s, i_{s+1}, \dots, i_n}^{l_1, \dots, l_s}$ can be chosen to be completely symmetric in the indices $\sigma, \nu_1, \dots, \nu_s$ and completely antisymmetric in the indices l_1, \dots, l_s .

9 Conclusions

We have succeeded to give a fairly complete presentation of the most important results connected with the existence of the variational sequences. Many of the proofs have been essentially simplified using Fock space techniques. We also have been able to give very explicit expression for the most general form of a variationally trivial Lagrangian and the generic expression of a locally variational system of partial differential equations, completing in this way known results from the literature.

Some criticism of the approach to the Lagrangian formalism accepted in this paper is necessary. We think that the weak point of this approach, from the physical point of view, is in fact, the definition of a differential equation. The definition of such an object as a special type of differential form leads at the following problem. Let T_σ, T'_σ be the components of two differential equations in a given chart. It is possible (and examples can be provided) that one can arrange such that: (1) the hyper-surface $T_\sigma = 0 \quad \sigma = 1, \dots, m$ coincides with the hyper-surface $T'_\sigma = 0 \quad \sigma = 1, \dots, m$ (in this way the two sets of functions are describing in fact the same set of physical solutions of the equation of motion) and (2) T_σ are locally variational and T'_σ are not locally variational. So, in a certain sense, the property of being locally variational is not intrinsically defined. (See on this point also [6]). In this sense, the most reasonable definition (from the physical point of view) of a differential equation would be certain hyper-surfaces in the jet bundle extension with some regularity properties (guaranteeing a well posed Cauchy problem). The problem would be to attach in an intrinsic way the property of being locally variational to such a hyper-surface.

Another interesting and open problem is to find, if possible, a physical meaning for the elements of the variational sequence of index $q \geq n + 2$ and eventually some representatives by forms.

One would also be interested to see to what extent the results of this approach to the Lagrangian formalism can be extended to the case when Y is not a fibre bundle over some "space-time" manifold X (the typical case being a relativistic system with Y the Minkowski space). Although it is clear that the line of argument from this paper depends essentially on the existence of the fibre bundle structure, some steps in this direction exists in the literature [9], [13]; the proper substitute for jet extensions of a fibre bundle are the higher order Grassmann bundles.

Finally, there exists some physical interest to extend this formalism to the situation when anticommuting variables are present. This case appears when one is studying, for instance, BRST-type symmetries.

10 Appendix

In this appendix we give the basic definitions of the Fock space concepts we have used in this paper and provide a fairly simple proof of the so-called *trace decomposition theorem* [24]. One can simplify somehow all the proofs in this paper if one uses this more refined decomposition of tensors; however this simplification is rather modest.

10.1 Fock Space Notions

To avoid unnecessary complications we consider \mathcal{H} to be a finite dimensional real Hilbert space. Then the *associated Fock space* is, by definition:

$$\mathcal{F}(\mathcal{H}) \equiv \oplus_{n=0}^{\infty} \mathcal{H}_n$$

where

$$\mathcal{H}_n \equiv \mathcal{H}^{\otimes n}, \quad n > 0, \quad \mathcal{H}_0 \equiv \mathbb{R}.$$

The Hilbert space $\mathcal{H} \sim \mathcal{H}_1$ is called the *one-particle space* and the element $(1, 0, \dots) \in \mathcal{F}(\mathcal{H})$ is called the *vacuum*.

We can introduce the symmetrization and the antisymmetrization operators in \mathcal{H}_n according to

$$\mathcal{S}_n^{\pm} \phi_1 \otimes \dots \otimes \phi_n \equiv \frac{1}{n!} \sum_{P \in \mathcal{P}_n} \epsilon_{\pm}(P) \phi_{P(1)} \otimes \dots \otimes \phi_{P(n)} \quad (10.1)$$

where

$$\epsilon_+(P) \equiv 1, \quad \epsilon_-(P) \equiv (-1)^{|P|}, \quad \forall P \in \mathcal{P}_n. \quad (10.2)$$

Here \mathcal{P}_n is the permutation group of the numbers $1, 2, \dots, n$ and $|P|$ is the signature of the permutation P . One can prove easily that these operators are in fact orthogonal projectors. We also define the following projector operators acting in the Fock space:

$$\mathcal{S}^{\pm} \equiv \oplus_{n=0}^{\infty} \mathcal{S}_n^{\pm}. \quad (10.3)$$

Next, one defines the following subspaces of \mathcal{H}_n :

$$\mathcal{H}_n^{\pm} \equiv \mathcal{S}_n^{\pm} \mathcal{H}_n \quad (10.4)$$

and of $\mathcal{F}(\mathcal{H})$:

$$\mathcal{F}^{\pm}(\mathcal{H}) \equiv \mathcal{S}^{\pm} \mathcal{F}(\mathcal{H}) = \oplus_{n=0}^{\infty} \mathcal{S}_n^{\pm} \mathcal{H}_n. \quad (10.5)$$

The subspace $\mathcal{F}^+(\mathcal{H})$ is called the *bosonic (or symmetric) Fock space* for the sign $+$ and the *fermionic (or antisymmetric) Fock space* for the sign $-$.

One can define in $\mathcal{F}(\mathcal{H})$ the so-called *particle number operators* according to

$$N \phi_1 \otimes \dots \otimes \phi_n \equiv n \phi_1 \otimes \dots \otimes \phi_n. \quad (10.6)$$

It is clear that these operators can be restricted to the bosonic and to the fermionic Fock spaces.

To simplify the presentation according to our specific needs, we consider an orthonormal basis in the $\mathcal{H} : e_1, \dots, e_k$, $k \equiv \dim(\mathcal{H})$. Then every element of the Fock space $\mathcal{F}(\mathcal{H})$ can be represented as a collection

$$(f, f^i, \dots, f^{i_1, \dots, i_n}, \dots)$$

where f^{i_1, \dots, i_n} are the elements of a (real) tensor of degree n .

The operators \mathcal{S}^\pm are represented by the following formulæ:

$$(\mathcal{S}^\pm f)^{i_1, \dots, i_n} \equiv \frac{1}{n!} \sum_{P \in \mathcal{P}_n} \epsilon_\pm(P) f^{i_{P(1)}, \dots, i_{P(n)}}, \quad \forall n > 0. \quad (10.7)$$

Sometimes it is convenient to indicate explicitly the indices affected by the operation of symmetrization, or antisymmetrization, by writing the preceding formula as follows:

$$\mathcal{S}_{i_1, \dots, i_n}^\pm f^{i_1, \dots, i_n} \equiv \frac{1}{n!} \sum_{P \in \mathcal{P}_n} \epsilon_\pm(P) f^{i_{P(1)}, \dots, i_{P(n)}}, \quad \forall n > 0. \quad (10.8)$$

This notation is important because we have the following formulæ which are used many times in the proofs:

$$\mathcal{S}_I^\pm \mathcal{S}_J^\pm = \mathcal{S}_I^\pm, \quad \forall J \subset I. \quad (10.9)$$

The bosonic (resp. fermionic) Fock space is formed by symmetric (resp. antisymmetric) tensors.

We are ready to introduce now the so-called *creation* and *annihilation* operators. We will not use the usual convention from quantum mechanics textbooks, but another one more suitable for our purposes. We also use consistently Bourbaki conventions:

$$\sum_{\emptyset} \equiv 0, \quad \prod_{\emptyset} \equiv 1.$$

Consider first the bosonic case. We define in the bosonic Fock space the *creation* and *annihilation* operators b_l^* , b^l , $l = 1, \dots, n$ by:

$$(b_l^* f)^{i_1, \dots, i_n} \equiv \frac{1}{n} \sum_{p=1}^n \delta_{l^p}^{i_p} f^{i_1, \dots, \hat{i}_p, \dots, i_n} = \mathcal{S}_{i_1, \dots, i_n}^+ \delta_{l^1}^{i_1} f^{i_2, \dots, i_n}, \quad l = 1, \dots, k, \quad \forall n \geq 0 \quad (10.10)$$

(where we prefer to specify explicitly the indices on which the operation of symmetrization is performed; evidently, for $n = 0$ the right hand side must be considered 0) and respectively by:

$$(b^l f)^{i_1, \dots, i_n} \equiv (n+1) f^{l, i_1, \dots, i_n}, \quad \forall n \geq 0. \quad (10.11)$$

It is easy to see that, in fact, b_l^* is the adjoint of b^l . The essential property of these operators is contained in the so-called *canonical commutation relations (CCR)*:

$$[b^l, b^m] = 0, \quad [b_l^*, b_m^*] = 0, \quad [b^l, b_m^*] = \delta_m^l, \quad \forall l, m = 1, \dots, k \quad (10.12)$$

where by $[\cdot, \cdot]$ we are designating, obviously, the commutator of the two entries.

We now consider the fermionic case. The definitions are in complete analogy with the previous ones. We define in the fermionic Fock space the *creation* and *annihilation* operators a_l^* , a^l , $l = 1, \dots, n$ by:

$$(a_l^* f)^{i_1, \dots, i_n} \equiv \frac{1}{n} \sum_{p=1}^n (-1)^{p-1} \delta_l^{i_p} f^{i_1, \dots, \hat{i}_p, \dots, i_n} = \mathcal{S}_{i_1, \dots, i_n}^- \delta_l^{i_1} f^{i_2, \dots, i_n}, \quad l = 1, \dots, k, \quad \forall n \geq 0 \quad (10.13)$$

(where again we specify explicitly the indices on which the operation of antisymmetrization is performed; for $n = 0$ the right hand side must be considered 0) and respectively by:

$$(a^l f)^{i_1, \dots, i_n} \equiv (n+1) f^{i_1, \dots, i_n, l}, \quad \forall n \geq 0. \quad (10.14)$$

Again one sees that a_l^* is the adjoint of a^l . We have analogously to (10.12) the *canonical anticommutation relations (CAR)*:

$$\{a^l, a^m\} = 0, \quad \{a_l^*, a_m^*\} = 0, \quad \{a^l, a_m^*\} = \delta_m^l, \quad \forall l, m = 1, \dots, k \quad (10.15)$$

where by $\{\cdot, \cdot\}$ we are designating the anticommutator of the two entries.

Remark 10.0.3 *It is not very complicated to find the abstract definitions for the creation and the annihilation operators, i.e. basis independent definitions, but we will not need them here. It is also noteworthy that the whole formalism works for any Hilbert space, even infinite dimensional, defined over an arbitrary commutative division field.*

It is obvious how to extend the definitions of the particle number operator and of the creation and annihilation operators to the more general case of the Hilbert space \mathcal{H}_s appearing in the proof of theorem 3.3. In particular, we have more particle number operators: N_f corresponding to the fermionic degrees of freedom and N_α , $\alpha = 1, \dots, s$ corresponding to the bosonic degrees of freedom and giving k (respectively r_α) when applied on $\mathcal{H}_{k, r_1, \dots, r_s}$.

In this case, if we define the operators B_α , $\alpha = 1, \dots, s$ according to (3.24) then it is easy to establish the following anticommutation relations:

$$\{B_\alpha, B_\alpha^*\} = N_f - N_\alpha + n\mathbf{1}, \quad \alpha = 1, \dots, s \quad (10.16)$$

and

$$\{B_\alpha, B_\beta\} = 0, \quad \alpha, \beta = 1, \dots, s; \quad (10.17)$$

in particular we have:

$$B_\alpha^2 = 0, \quad \alpha = 1, \dots, s. \quad (10.18)$$

We also have the commutation relations:

$$[B_\alpha, N_f - N_\alpha + n\mathbf{1}] = 0, \quad \alpha = 1, \dots, s. \quad (10.19)$$

10.2 The Trace Decomposition Formula

A tensor of the form $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ is called *traceless* iff it verifies:

$$A_{\sigma_1, \dots, \sigma_s, j, i_{s+1}, \dots, i_q}^{\{j, p_1, \dots, p_l\}, I_2, \dots, I_s} = 0 \quad (10.20)$$

for all indices left free. If we use compact tensor notations, this can be written as follows:

$$B_1^* A_{\sigma_1, \dots, \sigma_s} = 0. \quad (10.21)$$

Because of the symmetry properties (3.18) we have in fact:

$$B_\alpha^* A_{\sigma_1, \dots, \sigma_s} = 0 \quad (\alpha = 1, \dots, s). \quad (10.22)$$

Then we have the following result

Lemma 10.1 *Let $X \in \mathcal{H}_{p, r_1, \dots, r_s}$ ($r_1, \dots, r_s \in \mathbb{N}^*$, $s \leq n - p$). Then the following decomposition formula is valid:*

$$X = X_0 + \sum_{\alpha=1}^k B_\alpha X_\alpha \quad (10.23)$$

where the tensor X_0 is traceless and it is **uniquely** determined by X .

Proof:

It follows elementary from the relations at the end of the preceding subsection that the operators

$$P_\alpha \equiv (N_f - N_\alpha + n\mathbf{1})^{-1} B_\alpha B_\alpha^*, \quad Q_\alpha \equiv (N_f - N_\alpha + n\mathbf{1})^{-1} B_\alpha^* B_\alpha, \quad \alpha = 1, \dots, s \quad (10.24)$$

are orthogonal projectors and we have

$$P_\alpha + Q_\alpha = 1, \quad \alpha = 1, \dots, s. \quad (10.25)$$

We consider now the corresponding subspaces

$$V_\alpha \equiv \text{Span}(P_\alpha \mathcal{H}_s), \quad \alpha = 1, \dots, s. \quad (10.26)$$

It is well known that the set of all linear subspaces in a finite dimensional Hilbert space is a orthomodular lattice if the order relation $<$ is given by the inclusion of subspaces \subset and the orthogonalization operator is constructed using the scalar product [28]. It follows that one has the following formulæ for the *infimum* and *supremum* operations:

$$\bigwedge_{i \in I} V_i \equiv \bigcap_{i \in I} V_i, \quad \bigvee_{i \in I} V_i \equiv \sum_{i \in I} V_i. \quad (10.27)$$

The property above can be transported for the set of orthogonal projectors acting in this Hilbert space. In this new representation, the orthogonalization operation looks very simple:

$$P^\perp \equiv \mathbf{1} - P. \quad (10.28)$$

Now we decompose the generic element X as follows:

$$X = X_0 + X_1 \quad (10.29)$$

where

$$X_0 \equiv (\wedge Q_\alpha) X, \quad X_1 \equiv X - X_0. \quad (10.30)$$

One immediately establishes that X_0 is traceless and that

$$X_1 = (\wedge Q_\alpha)^\perp X = \vee Q_\alpha^\perp X = \vee P_\alpha X$$

so X_1 has the form:

$$X_1 = \sum_{\alpha} B_{\alpha} X_{\alpha}.$$

This finishes the proof. \blacksquare

Remark 10.1.1 *One can prove that the lattice property is valid in a more general case of linear spaces of Hilbertian type [28]).*

Remark 10.1.2 *A more general trace decomposition formula appears in [24] in the sense that one can give up the hypothesis that the Hilbert space is of Fock type.*

As a consequence we have

Theorem 10.2 *Let $\rho \in \Omega_q^r$ and let (V, ψ) be a chart on Y . Then the form ρ admits in the chart (V^r, ψ^r) the following **unique** decomposition:*

$$\rho = \rho_0 + \rho' \quad (10.31)$$

where $\rho_0 \in \Omega_{q(c)}^r$ is a contact form, and ρ' has the expression

$$\rho' = \sum_{s=0}^q \frac{1}{s!(q-s)!} \sum_{|I_1|, \dots, |I_s|=r} A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} dy_{I_1}^{\sigma_1} \wedge \dots \wedge dy_{I_s}^{\sigma_s} \wedge dx^{i_{s+1}} \wedge \dots \wedge dx^{i_q}, \quad (10.32)$$

where the tensors $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ verify the antisymmetry property (3.18) are traceless.

Proof:

We use the standard decomposition given by (3.15) - (3.17) and we refine it.

We apply the preceding lemma to the tensors $A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s}$ appearing in the expression of ρ' (see (3.17)). So, this tensor splits into a δ -contribution and a traceless contribution. The first contribution leads to terms containing at least a factor $d\omega_J^\sigma$ $|J| = r-1$ so it is a contact form and can be combined with ρ_0 ; the second contribution is traceless. This means that we have the desired decomposition. The uniqueness assertion is easy to prove. \blacksquare

Remark 10.2.1 *It immediately follows that the equivalence classes $\Omega_q^r(V) / \Omega_{q(c)}^r(V)$ ($q = 1, \dots, n$) (for any chart V on Y) are indexed by sets of traceless tensors*

$$A_{\sigma_1, \dots, \sigma_s, i_{s+1}, \dots, i_q}^{I_1, \dots, I_s} \quad (|I_1| = \dots = |I_s| = r, \quad s = 0, \dots, q).$$

For the proof of the isomorphism $\Omega_q^r(V) / \Omega_{q(c)}^r(V) \cong \mathcal{J}_q^{r+1}(V)$ ($q = 1, \dots, n$) see the end of [23].

References

- [1] S. J. Aldersley, “*Higher Euler Operators and some of their Applications*”, Journ. Math. Phys. **20** (1979) 522-531
- [2] I. M. Anderson, “*The Variational Bicomplex*”, Utah State Univ. preprint, 1989, (Academic Press, Boston, to appear)
- [3] I. M. Anderson, T. Duchamp, “*On the Existence of Global Variational Principles*”, American Journ. Math. **102** (1980) 781-868
- [4] I. M. Anderson, J. Pohjanpetlo, “*Variational Principles for Differential Equations with Symmetry and Conservation Laws. I. Second Order Scalar Equations*”, Math. Ann. **299** (1994) 191-222
- [5] I. M. Anderson, J. Pohjanpetlo, “*Symmetries, Conservation Laws and Variational Principles for Vector Field Theories*”, preprint, 1995
- [6] I. M. Anderson, G. Thompson, “*The Inverse Problem of the Calculus of Variation for Ordinary Differential Equations*”, Memoirs of AMS, vol. **98**, no. **473**, 1992
- [7] J. M. Ball, J. C. Currie, P. J. Olver, “*Null Lagrangians, Weak Continuity, and Variational Problems of Arbitrary Order*”, Journ. Functional Anal. **41** (1981) 135-174
- [8] D. G. Edelen, “*The Null Set of the Euler-Lagrange Operator*”, Arch. Rat. Mech. Anal. **11** (1962) 117-121
- [9] D. R. Grigore, “*A Generalized Lagrangian Formalism in Particle Mechanics and Classical Field Theory*”, Fortschr. der Phys. **41** (1993) 569-61
- [10] D. R. Grigore, “*Variational Equations and Symmetries in the Lagrangian Formalism*”, J. Phys. **A 28** (1995) 2921-2937
- [11] D. R. Grigore, “*Variational Equations and Symmetries in the Lagrangian Formalism; Arbitrary Vector Fields*”, hep-th/9512176, to appear in Fortschr. der Phys.
- [12] D. R. Grigore, “*Trivial Second-Order Lagrangians in Classical Field Theory*”, hep-th/9609199, submitted for publication
- [13] D. R. Grigore, D. Krupka, “*Invariants of Velocities and Higher Order Grassmann Bundles*”, dg-ga/9708013, to appear in Journ. Geom. Phys.
- [14] D. R. Grigore, O. T. Popp, “*On the Lagrange-Souriau Form in Classical Field Theory*”, to appear in Mathematica Bohemica
- [15] P. J. Olver, “*Hyperjacobians, Determinant Ideals and the Weak Solutions to Variational Problems*”, Proc. Roy. Soc. Edinburgh **95A** (1983) 317-340

- [16] D. Krupka, “*A Geometric Theory of Ordinary First Order Variational Problems in Fibered Manifolds. I. Critical Sections*”, Journ. Math. Analysis Appl. **49** (1975) 180-206
- [17] D. Krupka, “*On the Local Structure of the Euler-Lagrange Mapping of the Calculus of Variations*”, in Proceedings of the Conference on Differential Geometry and its Applications, (September, 1981), published by Univ. Karlova, pp. 181-188
- [18] D. Krupka, “*Lepagean Forms in Higher Order Variational Theory*”, in Proceedings of the IUTAM-ISIMM Symposium on “Modern Developments in Analytical Mechanics”, (Turin, 1982), Atti della Accademia delle Scienze di Torino, Suppl. al Vol. **117** (1983) 198-238
- [19] D. Krupka, “*On the Uniqueness of Some Differential Invariants: $d, [,], \nabla$* ”, Czechoslovak Math. Journ. **34** (1984) 588-597
- [20] D. Krupka, “*Geometry of Lagrangian Structures.3*”, in Proceedings of the 14th Winter School on Abstract Analysis, SRNI, 1986, Suppl. ai Rendiconti del Circolo Matematico di Palermo, Serie II, no.14, 1987, pp. 187-224
- [21] D. Krupka, “*Variational Sequence on Finite Order Jet Spaces*”, in Proceedings of the Conference “Differential Geometry and its Applications”, Brno, August, 1989 (World Scientific: Singapore, 1990), pp. 236-254
- [22] D. Krupka, “*Topics in the Calculus of Variation: Finite Order Variational Sequences*”, in “Differential Geometry and its Applications”, proceedings Conf. Opava, 1992, (Opava: Open Univ. Press), pp. 437-495
- [23] D. Krupka, “*The Contact Ideal*”, Diff. Geometry and its Applications **5** (1995) 257-276
- [24] D. Krupka, “*The Trace Decomposition Problem*”, Beiträge zur Algebra und Geometrie **36** (1995) 303-315
- [25] H. Rund, “*Integral Formulæ Associated with the Euler-Lagrange Operator of Multiple Integral Problems in the Calculus of Variation*”, Æquationes Math. **11** (1974) 212-229
- [26] F. Takens, “*Symmetries, Conservation Laws and Variational Principles*”, in “Geometry and Topology”, proceedings of the III Latin American School of Mathematics (Rio de Janeiro, 1976), Lect. Notes in Math. **599**, (Berlin: Springer, 1977), J. Palais, M. de Carmo (eds.), pp. 581-604
- [27] W. M. Tulczyjew, “*The Euler-Lagrange Resolution*”, in Int. Coll. on Differential Geometrical Methods in Math. Phys. (Aix-en-Provence, 1979), Lect. Notes. in Math. **836** (Berlin: Springer), pp. 22-48
- [28] V. S. Varadarajan, “*Geometry of Quantum Theory*”, van Nostrand Co., Princeton, 1968